arXiv:1110.3192v1 [math.DS] 14 Oct 2011

INTERSECTIONS OF HOMOGENEOUS CANTOR SETS AND BETA-EXPANSIONS

DERONG KONG, WENXIA LI, AND MICHEL DEKKING

Abstract. Let $\Gamma_{\beta,N}$ be the N-part homogeneous Cantor set with $\beta \in (1/(2N-1), 1/N)$. Any string $(j_\ell)_{\ell=1}^{\infty}$ with $j_\ell \in \{0, \pm 1, \ldots, \pm(N-1)\}$ such that $t = \sum_{\ell=1}^{\infty} j_\ell \beta^{\ell-1}(1-\beta)/(N-1)$ is called a code of t. Let $\mathcal{U}_{\beta,\pm N}$ be the set of $t \in [-1,1]$ having a unique code, and let $\mathcal{S}_{\beta,\pm N}$ be the set of $t \in \mathcal{U}_{\beta,\pm N}$ which make the intersection $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a self-similar set. We characterize the set $\mathcal{U}_{\beta,\pm N}$ in a geometrical and algebraical way, and give a sufficient and necessary condition for $t \in \mathcal{S}_{\beta,\pm N}$. Using techniques from beta-expansions, we show that there is a critical point $\beta_c \in (1/(2N-1), 1/N)$, which is a transcendental number, such that $\mathcal{U}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \beta_c)$, and contains countably infinite many elements if $\beta \in (\beta_c, 1/N)$. Moreover, there exists a second critical point $\alpha_c = \left[N+1-\sqrt{(N-1)(N+3)}\right]/2 \in (1/(2N-1), \beta_c)$ such that $\mathcal{S}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \alpha_c)$, and contains countably infinite many elements if $\beta \in [\alpha_c, 1/N)$.

Keywords: Homogeneous Cantor set; self-similarity; iterated function system; critical point; beta-expansion; Thue-Morse sequence.

MSC: 28A80, 28A78

1. Introduction

Let $\{f_i(x) = r_i x + b_i\}_{i=1}^{p}$ be a family of functions on $\mathbb{R}$ with $0 < |r_i| < 1$. It is well known (cf. [5]) that there exists a unique nonempty compact set $\Gamma \subseteq \mathbb{R}$ such that

$$\Gamma = \bigcup_{i=1}^{p} f_i(\Gamma).$$

In this case, Γ is called the *self-similar set* generated by the *iterated function system* (IFS) $\{f_i(\cdot)\}_{i=1}^{p}$.

We will be interested in the self-similar set $\Gamma_{\beta,\Omega}$ generated by an IFS $\{\phi_d(\cdot) : d \in \Omega\}$, where Ω is a finite set of integers, and

$$\phi_d(x) = \beta x + d(1-\beta)/(N-1), \quad x \in \mathbb{R}$$

Date: October 17, 2011.

The second author is supported by the National Natural Science Foundation of China no 10971069, Shanghai Leading Academic Discipline Project no B407 and Shanghai Education Committee Project no 11ZZ41.

for some $N \geq 2$ and $\beta \in (0, 1/N)$. It is well known that one can establish a surjective map $\pi_\Omega : \Omega^\infty \to \Gamma_{\beta,\Omega}$ by letting

$$\pi_\Omega(J) = \sum_{\ell=1}^{\infty} \frac{j_\ell \beta^{\ell-1}(1-\beta)}{N-1} \tag{1}$$

for $J = (j_\ell)_{\ell=1}^\infty \in \Omega^\infty$. The infinite string J is called an Ω-*code* of $\pi_\Omega(J)$. Note that an element $x \in \Gamma_{\beta,\Omega}$ may have multiple Ω-codes. These Ω-codes are closely related to the classical beta-expansions (cf. [4, 7, 12, 17, 18, 19, 20]). A sequence $(s_\ell)_{\ell=1}^\infty \in \Omega^\infty$ is called a *β-expansion of x with digit set* Ω if we can write

$$x = \sum_{\ell=1}^{\infty} s_\ell \beta^\ell, \quad s_\ell \in \Omega.$$

Let $\Omega_N := \{0, 1, \ldots, N-1\}$. We simplify the notation Γ_{β,Ω_N} to $\Gamma_{\beta,N}$, so this set satisfies

$$\Gamma_{\beta,N} = \bigcup_{d \in \Omega_N} \phi_d(\Gamma_{\beta,N}).$$

The set $\Gamma_{\beta,N}$ is called the *N-part homogeneous Cantor set*. Thus $\Gamma_{1/3,2}$ is the *classical middle-third Cantor set* and $\Gamma_{\beta,2}$ is the middle-α Cantor set with $\alpha = 1 - 2\beta$.

In terms of (1), let $\pi_N := \pi_{\Omega_N}$. Thus we can rewrite $\Gamma_{\beta,N}$ as

$$\Gamma_{\beta,N} = \pi_N\left(\Omega_N^\infty\right) = \left\{ \sum_{\ell=1}^{\infty} \frac{j_\ell \beta^{\ell-1}(1-\beta)}{N-1} : j_\ell \in \Omega_N \right\}. \tag{2}$$

We consider the intersection of $\Gamma_{\beta,N}$ with its translation by t. It is easy to check that

$$\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) \neq \emptyset \quad \text{if and only if} \quad t \in \Gamma_{\beta,N} - \Gamma_{\beta,N}.$$

Here we denote for a real number a, and sets $A, B \subseteq \mathbb{R}$, $aA := \{ax : x \in A\}$, $A + B := \{x + y : x \in A, y \in B\}$, and $A + a := A + \{a\}$.

It follows from Equation (2) that the difference set $\Gamma_{\beta,N} - \Gamma_{\beta,N}$ can be written as

$$\Gamma_{\beta,N} - \Gamma_{\beta,N} = \left\{ \sum_{k=1}^{\infty} \frac{t_\ell \beta^{\ell-1}(1-\beta)}{N-1} : t_\ell \in \Omega_{\pm N} \right\} = \pi_{\pm N}\left(\Omega_{\pm N}^\infty\right) = \Gamma_{\beta,\Omega_{\pm N}},$$

where $\Omega_{\pm N} := \Omega_N - \Omega_N = \{0, \pm 1, \ldots, \pm(N-1)\}$ and $\pi_{\pm N} := \pi_{\Omega_{\pm N}}$. Since $\Omega_{2N-1} = \{0, 1, \ldots, 2N-2\} = \Omega_{\pm N} + N - 1$, it is easy to see that $(t_\ell)_{\ell=1}^\infty$ is a $\Omega_{\pm N}$-code of $t \in \Gamma_{\beta,N} - \Gamma_{\beta,N}$ if and only if $(t_\ell + N - 1)_{\ell=1}^\infty$ is an β-expansion of $(t+1)\beta(N-1)/(1-\beta)$ with digit set Ω_{2N-1}. Thus some results and techniques from beta-expansions can be used to deal with the difference set $\Gamma_{\beta,N} - \Gamma_{\beta,N}$.

In the past two decades, intersections of Cantor sets have been studied by several authors (cf. [2, 8, 9, 10, 11, 13]). Recently, Deng et al. [3] gave a necessary and sufficient condition for $t \in [-1, 1]$ such that $\Gamma_{1/3,2} \cap (\Gamma_{1/3,2} + t)$ is a self-similar set. Their results were extended to the case $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ with $\beta \in (0, 1/(2N-1)]$ by Li et al. [15], and to the case $\Gamma_{\beta,2} \cap (\Gamma_{\beta,2} + t)$ with $\beta \in (1/3, 1/2)$ and t having a unique $\Omega_{\pm 2}$-code by Zou et al. [21].

In this paper we consider arbitrary $N \geq 2$, and $\beta \in (1/(2N-1), 1/N)$. Then Lebesgue a.a. $t \in \Gamma_{\beta,N} - \Gamma_{\beta,N} = [-1, 1]$ have a continuum of distinct $\Omega_{\pm N}$-codes. This gives the set

$\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a more complicated structure. We summarize the results in the following. In Section 2, an algebraical and geometrical description of the set

$$\mathcal{U}_{\beta,\pm N} := \big\{t \in [-1,1] : |\pi_{\pm N}^{-1}(t)| = 1\big\}$$

(i.e., the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code) is given in Theorem 2.2, where throughout the paper $|A|$ denotes the number of members in the set A. Section 3 is mainly devoted to investigating the self-similar structure of $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$. Let

$$\mathcal{S}_{\beta,\pm N} := \big\{t \in \mathcal{U}_{\beta,\pm N} : \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) \text{ is a self-similar set}\big\}.$$

Theorem 3.2 gives a sufficient and necessary condition for $t \in \mathcal{S}_{\beta,\pm N}$. In Section 4, we study the set $\mathcal{U}_{\beta,\pm N}$ for different $\beta \in (1/(2N-1), 1/N)$ culminating in Theorem 4.6. Using techniques from beta-expansions, we obtain a *critical point* $\beta_c \in (1/(2N-1), 1/N)$ such that $\mathcal{U}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \beta_c)$, and contains countably infinite many elements if $\beta \in (\beta_c, 1/N)$. We point out that the critical point β_c is a transcendental number which is related to the famous Thue-Morse sequence (cf. [12]). In Section 5 we find the second critical point $\alpha_c = [N + 1 - \sqrt{(N-1)(N+3)}\,]/2 \in (1/(2N-1), \beta_c)$ (see Theorem 5.1) such that $\mathcal{S}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \alpha_c)$, and contains countably infinite many elements if $\beta \in [\alpha_c, 1/N)$. In the following table, we give the critical points $\beta_c = \beta_c(N)$ and $\alpha_c = \alpha_c(N)$ calculated for different integers N by means of Mathematica.

N	2	3	4	5	6	7	8	9
$\beta_c \approx$	0.39433	0.27130	0.21004	0.17221	0.14625	0.12722	0.11265	0.10111
$\alpha_c \approx$	0.38197	0.26795	0.20871	0.17157	0.14590	0.12702	0.11252	0.10102

Thus for $\beta \in [\alpha_c, \beta_c)$, the set $\mathcal{U}_{\beta,\pm N}$ (the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code) has positive Hausdorff dimension, but only countably many $t \in \mathcal{U}_{\beta,\pm N}$ make the intersection $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a self-similar set.

2. GEOMETRICAL DESCRIPTION OF $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$

We say that the IFS $\{f_i(\cdot)\}_{i=1}^{p}$ satisfies the *open set condition* (OSC) if there exists a nonempty bounded open set $O \subseteq \mathbb{R}$ such that $O \supseteq \bigcup_{i=1}^{p} f_i(O)$, with a disjoint union on the right side. An IFS $\{f_i(\cdot)\}_{i=1}^{p}$ is said to satisfy the *strong separation condition* (SSC) if the union $\Gamma = \bigcup_{i=1}^{p} f_i(\Gamma)$ is disjoint.

When $\beta \in (0, 1/(2N-1))$ the IFS $\{\phi_d(\cdot) : d \in \Omega_{\pm N}\}$ satisfies the SSC, so each point in $\Gamma_{\beta,\Omega_{\pm N}}$ has a unique $\Omega_{\pm N}$-code. In case $\beta = 1/(2N-1)$, the IFS $\{\phi_d(\cdot) : d \in \Omega_{\pm N}\}$ fails to satisfy the SSC but satisfies the OSC, so each point has a unique $\Omega_{\pm N}$-code except for countably many points having two $\Omega_{\pm N}$-codes. However, for the case $\beta \in (1/(2N-1), 1/N)$ the IFS $\{\phi_d(\cdot) : d \in \Omega_{\pm N}\}$ fails to satisfy the OSC and $\Gamma_{\beta,\Omega_{\pm N}} = [-1,1]$. In this case, Lebesgue a.a. $t \in [-1,1]$ have a continuum of distinct $\Omega_{\pm N}$-codes (cf. [19]). This gives

$\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a more complicated structure, since it follows ([13]) that for $t \in \Gamma_{\beta,\Omega_{\pm N}}$

$$\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) = \bigcup_{\tilde{t}} \pi_N \left(\prod_{\ell=1}^{\infty} D_{\ell,\tilde{t}} \right) \tag{3}$$

where the union is taken over all $\Omega_{\pm N}$-codes of t, and for each code $\tilde{t} = (t_\ell)_{\ell=1}^{\infty} \in \Omega_{\pm N}^{\infty}$

$$D_{\ell,\tilde{t}} = \Omega_N \cap (\Omega_N + t_\ell) = \{0, 1, \ldots, N-1\} \cap (\{0, 1, \ldots, N-1\} + t_\ell).$$

Moreover, $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ has the following properties:

(P1) the union on the right side of (3) consists of pairwise disjoint sets;

(P2) for each $\Omega_{\pm N}$-code $\tilde{t} = (t_\ell)_{\ell=1}^{\infty}$ of t, we have

$$1 + t - \pi_N \left(\prod_{\ell=1}^{\infty} D_{\ell,\tilde{t}} \right) = \pi_N \left(\prod_{\ell=1}^{\infty} D_{\ell,\tilde{t}} \right),$$

i.e., $\pi_N(\prod_{\ell=1}^{\infty} D_{\ell,\tilde{t}})$ is centrally symmetric. Furthermore, $1 + t - \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) = \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$.

These properties can be obtained as follows. Let $(t_\ell)_{\ell=1}^{\infty}$ be a $\Omega_{\pm N}$-code of t and let $J = (j_\ell)_{\ell=1}^{\infty} \in \Omega_N^{\infty}$. If

$$\pi_N(J) = \sum_{\ell=1}^{\infty} \frac{j_\ell \beta^{\ell-1}(1-\beta)}{N-1} \in \pi_N \left(\prod_{\ell=1}^{\infty} \Omega_N \cap (\Omega_N + t_\ell) \right),$$

then $(j_\ell - t_\ell)_{\ell=1}^{\infty} \in \Omega_N^{\infty}$. Note that the IFS $\{\phi_d(\cdot) : d \in \Omega_N\}$ satisfies the SSC (since $\beta < 1/N$). This implies that each point $x \in \Gamma_{\beta,N}$ has a unique Ω_N-code. Thus $(j_\ell - t_\ell)_{\ell=1}^{\infty}$ is the unique Ω_N-code of $\pi_N(J) - t$, implying (P1). In addition, one can check that for each $\ell \geq 1$,

$$N - 1 + t_\ell - \Omega_N \cap (\Omega_N + t_\ell) = \Omega_N \cap (\Omega_N + t_\ell),$$

implying (P2).

Let Ω be a nonempty finite subset of $\mathbb{Z}$. Denote by ε the empty word and put $\Omega^0 = \{\varepsilon\}$. For $I \in \bigcup_{\ell=0}^{\infty} \Omega^\ell$ and $J \in \Omega^\infty \cup \bigcup_{\ell=0}^{\infty} \Omega^\ell$, let $IJ \in \Omega^\infty \cup \bigcup_{\ell=0}^{\infty} \Omega^\ell$ be the concatenation of I and J. So in particular $\varepsilon J = J$. For a nonnegative integer k and a finite string $I \in \bigcup_{\ell=1}^{\infty} \Omega^\ell$, let $I^k := \overbrace{I \ldots I}^{k}$ be the k times repeating of I and $I^\infty := III \cdots \in \Omega^\infty$ be the infinite repeating of I. In particular, $I^0 = \varepsilon$. For $J = (j_\ell)_{\ell=1}^{\infty} \in \Omega^\infty$ and $k \in \mathbb{N}$, let $J|_k = (j_\ell)_{\ell=1}^{k} \in \Omega^k$. We define the algebraic difference between two infinite strings $I = (i_\ell)_{\ell=1}^{\infty}, J = (j_\ell)_{\ell=1}^{\infty} \in \Omega^\infty$ by $I - J = (i_\ell - j_\ell)_{\ell=1}^{\infty}$, and for a positive integer k let $I|_k - J|_k = (I - J)|_k = (i_\ell - j_\ell)_{\ell=1}^{k}$.

Given $\beta \in (1/(2N-1), 1/N)$ and $t \in [-1, 1]$, for an integer $d \in \mathbb{Z}$, let

$$\psi_d(x) = \beta x + d(1-\beta)/(N-1) + t(1-\beta), \quad x \in \mathbb{R}.$$

Then

$$\Gamma_{\beta,N} + t = \bigcup_{d \in \Omega_N} \psi_d(\Gamma_{\beta,N} + t).$$

For $J=(j_\ell)_{\ell=1}^k\in\Omega_N^k$ with $k\in\mathbb{N}$, let $\psi_J:=\psi_{j_1}\circ\cdots\circ\psi_{j_k}$ (the same for ϕ_J). For a real number x, it is easy to see that $\psi_d(t+x)=\phi_d(x)+t$ for all $d\in\Omega_N$. Thus by induction we obtain

$$\psi_J(t+x)=\phi_J(x)+t\quad\text{for all } J\in\bigcup_{\ell=1}^\infty\Omega_N^\ell,\ x\in\mathbb{R}. \tag{4}$$

The sets $\Gamma_{\beta,N}$ and $\Gamma_{\beta,N}+t$ can be represented in a geometrical way as (cf. [5])

$$\Gamma_{\beta,N}=\bigcap_{k=1}^\infty\bigcup_{J\in\Omega_N^k}\phi_J([0,1])\quad\text{and}\quad\Gamma_{\beta,N}+t=\bigcap_{k=1}^\infty\bigcup_{J\in\Omega_N^k}\psi_J([t,1+t]).$$

We call $\phi_J([0,1]),\psi_J([t,1+t])$ with $J\in\Omega_N^k$ the *k-level components* of $\Gamma_{\beta,N}$ and $\Gamma_{\beta,N}+t$, respectively. The 1-level components of $\Gamma_{\beta,N}$ are $\phi_0([0,1]),\phi_1([0,1]),\ldots,\phi_{N-1}([0,1])$ of length β. All gaps between them have the same length $(1-\beta)/(N-1)-\beta$. The left endpoint of $\phi_0([0,1])$ is 0 and the right endpoint of $\phi_{N-1}([0,1])$ is 1. For a ℓ-level component $\phi_J([0,1]),J\in\Omega_N^\ell$, the $(\ell+1)$-level components $\phi_{J0}([0,1]),\phi_{J1}([0,1]),\ldots,\phi_{J(N-1)}([0,1])$ have the same length $\beta^{\ell+1}$ and all gaps (called $(\ell+1)$-level gaps) between them have the same length $\beta^\ell(1-\beta)/(N-1)-\beta^{\ell+1}$. The left endpoint of $\phi_{J0}([0,1])$ coincides with the left endpoint of $\phi_J([0,1])$ and the right endpoint of $\phi_{J(N-1)}([0,1])$ coincides with the right endpoint of $\phi_J([0,1])$. The requirement $\beta\in(1/(2N-1),1/N)$ implies the following simple properties:

(P3) the length of a k-level gap is less than the length of a k-level component, i.e.,

$$\beta^{k-1}(1-\beta)/(N-1)-\beta^k<\beta^k;$$

(P4) if $\phi_I([0,1])\cap\psi_J([t,t+1])\neq\emptyset$ for $I,J\in\Omega_N^k$ with $k\in\mathbb{N}$, then

$$\phi_I([0,1])\cap\psi_J([t,1+t])\cap\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)\neq\emptyset.$$



FIGURE 1. $N=3$, $\beta=0.28$, $t=0.19$. The 1-level components of $\Gamma_{\beta,N}$ are $\phi_0([0,1])$, $\phi_1([0,1])$ and $\phi_2([0,1])$. The 1-level components of $\Gamma_{\beta,N}+t$ are $\psi_0([t,1+t])$, $\psi_1([t,1+t])$ and $\psi_2([t,1+t])$. Here $\mathcal{N}_t(0)=\{\psi_0([t,1+t])\}$, $\mathcal{N}_t(1)=\{\psi_0([t,1+t]),\ \psi_1([t,1+t])\}$ and $\mathcal{N}_t(2)=\{\psi_1([t,1+t]),\ \psi_2([t,1+t])\}$.

For $J\in\Omega_N^k$ with $k\in\mathbb{N}$, the *neighborhood* of $\phi_J([0,1])$ with respect to the $k-$level components of $\Gamma_{\beta,N}+t$ is defined as (see Figure 1)

$$\mathcal{N}_t(J):=\Big\{\psi_I([t,1+t]):I\in\Omega_N^k,\ \phi_J([0,1])\cap\psi_I([t,1+t])\neq\emptyset\Big\}.$$

The set $\mathcal{N}_t(J)$ may be empty and $|\mathcal{N}_t(J)| \in \{0,1,2\}$. For $k \geq 1$ let

$$\Lambda_k := \Big\{ J \in \Omega_N^k : |\mathcal{N}_t(J)| \geq 1 \Big\} \quad \text{and} \quad \Lambda := \Big\{ J \in \Omega_N^\infty : J|_k \in \Lambda_k \text{ for all } k \in \mathbb{N} \Big\}.$$

Then $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ can be rewritten in a geometrical way as

$$\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) = \pi_N(\Lambda) = \bigcap_{k=1}^{\infty} \bigcup_{J \in \Lambda_k} \phi_J([0,1]).$$

A set $D \subseteq \Omega_N$ is said to be *consecutive* if $D = \Omega_N \cap (\Omega_N + d)$ for some $d \in \Omega_{\pm N}$.

Proposition 2.1. *Given $N \geq 2$ and $\beta \in (1/(2N-1), 1/N)$, let $t \in [-1,1]$. If $|\mathcal{N}_t(J)| \leq 1$ for all $J \in \bigcup_{\ell=1}^\infty \Omega_N^\ell$, then*

$$\Lambda = \prod_{\ell=1}^{\infty} D_\ell$$

with each D_ℓ consecutive.

Proof. The condition $\beta \in (1/(2N-1), 1/N)$ implies (P3), i.e., all gaps between the intervals $\phi_d([0,1]), d \in \Omega_N$ have the same length strictly less than β, the length of $\phi_d([0,1])$ (see Figure 1). Thus since $t \in [-1,1]$, either $|\mathcal{N}_t(0)| = 1$ or $|\mathcal{N}_t(N-1)| = 1$, which implies that

$$D_1 := \Big\{ d \in \Omega_N : |\mathcal{N}_t(d)| = 1 \Big\} \neq \emptyset.$$

It follows from $|\mathcal{N}_t(d)| \leq 1$ for all $d \in \Omega_N$ that D_1 is consecutive and $\Lambda_1 = D_1$.

Now for $k \in \mathbb{N}$ let the consecutive sets $D_1, \ldots, D_k$ be chosen such that $\Lambda_k = \prod_{\ell=1}^k D_\ell$. Fix a $J \in \Lambda_k$ and take

$$D_{k+1} := \Big\{ d \in \Omega_N : |\mathcal{N}_t(Jd)| = 1 \Big\}.$$

Then D_{k+1} is nonempty by (P3), and is consecutive by the same argument as above. Note that D_{k+1} is independent of the choice of $J \in \Lambda_k$. Thus $\Lambda_{k+1} = \prod_{\ell=1}^{k+1} D_\ell$ which implies $\Lambda = \prod_{\ell=1}^\infty D_\ell$ by induction. □

The following theorem characterizes the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code from a geometrical and an algebraical aspect.

Theorem 2.2. *Given $N \geq 2$ and $\beta \in (1/(2N-1), 1/N)$, let $\mathcal{U}_{\beta,\pm N}$ be the set of $t \in [-1,1]$ which have a unique $\Omega_{\pm N}$-code. Then the following conditions are equivalent.*

(A) $t \in \mathcal{U}_{\beta,\pm N}$;

(B) $|\mathcal{N}_t(J)| \leq 1$ *for all* $J \in \bigcup_{\ell=1}^\infty \Omega_N^\ell$;

(C) t *has a* $\Omega_{\pm N}$*-code* $(t_\ell)_{\ell=1}^\infty$ *such that for all* $k \geq 1$

$$\begin{cases} \sum_{\ell=1}^\infty t_{k+\ell}\beta^\ell < \frac{1-N\beta}{1-\beta}, & \textit{if} \quad t_k < N-1 \\ \sum_{\ell=1}^\infty t_{k+\ell}\beta^\ell > -\frac{1-N\beta}{1-\beta}, & \textit{if} \quad t_k > 1-N. \end{cases} \tag{5}$$

Proof. (A) ⇒ (B). Suppose that $|\mathcal{N}_t(J)| = 2$ for some $J = (j_\ell)_{\ell=1}^k \in \Omega_N^k$ with $k \geq 1$. Then either $|\mathcal{N}_t(J|_{k-1}0)| = 2$ or $|\mathcal{N}_t(J|_{k-1}(N-1))| = 2$. Without loss of generality, let $|\mathcal{N}_t(J|_{k-1}0)| = 2$. Then there exists $d \in \Omega_N$ such that $|\mathcal{N}_t(J|_{k-1}d)| = 1$ by the geometric structure of $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ (see Figure 2).



FIGURE 2. $N = 3$. Here $J' = J|_{k-1}1$, $J'' = J|_{k-1}2$ and $\mathcal{N}_t(J') \cap \mathcal{N}_t(J'') = \{\psi_I([t, 1+t])\}$.

Let $J' = J|_{k-1}(d-1)$ and $J'' = J|_{k-1}d$. Then

$$\mathcal{N}_t(J') \cap \mathcal{N}_t(J'') = \big\{\psi_I([t, 1+t])\big\}$$

for some $I = i_1 i_2 \cdots i_{k-1}(N-1) \in \Omega_N^k$. By (P4) we can pick

$$x \in \phi_{J'}([0,1]) \cap \psi_I([t,1+t]) \cap \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$$

and

$$y \in \phi_{J''}([0,1]) \cap \psi_I([t,1+t]) \cap \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t).$$

Let $(x_\ell)_{\ell=1}^\infty$ and $(y_\ell)_{\ell=1}^\infty$ be the unique Ω_N-code of x and y, respectively. Then $x_k = d-1$ and $y_k = d$. On the other hand, $x-t, y-t \in \Gamma_{\beta,N}$ and by $(x_\ell^*)_{\ell=1}^\infty, (y_\ell^*)_{\ell=1}^\infty$ we denote their unique Ω_N-code, respectively. It follows from (4) that

$$x \in \psi_I([t,1+t]) = \phi_I([0,1]) + t \text{ and } y \in \psi_I([t,1+t]) = \phi_I([0,1]) + t,$$

which imply $x-t, y-t \in \phi_I([0,1])$. Thus $x_k^* = y_k^* = N-1$. Hence $t = x-(x-t) = y-(y-t)$ has two distinct $\Omega_{\pm N}$-codes: $(x_\ell - x_\ell^*)_{\ell=1}^\infty$ and $(y_\ell - y_\ell^*)_{\ell=1}^\infty$.

(B) $\Rightarrow$ (A). By Proposition 2.1, we have $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) = \pi_N(\prod_{\ell=1}^\infty D_\ell)$ with D_ℓ consecutive. Thus, it follows from (3) that t has a unique $\Omega_{\pm N}$-code $(t_\ell)_{\ell=1}^\infty$ with each t_ℓ determined by $D_\ell = \Omega_N \cap (\Omega_N + t_\ell)$.

(B) $\Rightarrow$ (C). It follows from Proposition 2.1 that $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) = \pi_N(\prod_{\ell=1}^\infty D_\ell)$ with each D_ℓ consecutive. Take $J = (j_\ell)_{\ell=1}^\infty \in \prod_{\ell=1}^\infty D_\ell$. Then $\pi_N(J) \in \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$. Let $J^* = (j_\ell^*)_{\ell=1}^\infty$ be the unique Ω_N-code of $\pi_N(J) - t \in \Gamma_{\beta,N}$. Thus it follows by (4) that for each $k \geq 1$

$$\pi_N(J) \in \phi_{J|_k}([0,1]) \cap (\phi_{J^*|_k}([0,1]) + t) = \phi_{J|_k}([0,1]) \cap \psi_{J^*|_k}([t, 1+t]),$$

and

$$J - J^* = (j_\ell - j_\ell^*)_{\ell=1}^\infty = (t_\ell)_{\ell=1}^\infty$$

is the unique $\Omega_{\pm N}$-code of t (the uniqueness is given by $(B) \Rightarrow (A)$). We shall prove $(t_\ell)_{\ell=1}^\infty$ satisfies (5) in the following.

Case I. $t_k \neq \pm(N-1)$.

In this case, $(j_k, j_k^*) \notin \{(N-1, 0), (0, N-1)\}$. This together with the requirements in (B) imply that the distance between the left endpoints of $\phi_{J|_k}([0,1])$ and $\psi_{J^*|_k}([t, t+1])$ must be less than the length of the k-th gap (see Figure 3), i.e., $|\psi_{J^*|_k}(t) - \phi_{J|_k}(0)| < \beta^{k-1}(1-\beta)/(N-1) - \beta^k$.



FIGURE 3. $N=3$. Here $\phi_{J|_k}(0)$ is the left endpoint of the k-level component $\phi_{J|_k}([0,1])$ of $\Gamma_{\beta,N}$, and $\psi_{J^*|_k}(t)$ is the left endpoint of k-level component $\phi_{J^*|_k}([t,1+t])$ of $\Gamma_{\beta,N}+t$.

Thus (5) follows by the following computation.

$$\begin{aligned}
\left|\sum_{\ell=1}^{\infty}\frac{t_{k+\ell}\beta^{\ell-1}(1-\beta)}{N-1}\right| &= \beta^{-k}\left|\sum_{\ell=k+1}^{\infty}\frac{t_\ell\beta^{\ell-1}(1-\beta)}{N-1}\right| = \beta^{-k}\left|t-\sum_{\ell=1}^{k}\frac{t_\ell\beta^{\ell-1}(1-\beta)}{N-1}\right| \\
&= \beta^{-k}\left|t-\left(\sum_{\ell=1}^{k}\frac{j_\ell\beta^{\ell-1}(1-\beta)}{N-1}-\sum_{\ell=1}^{k}\frac{j^*_\ell\beta^{\ell-1}(1-\beta)}{N-1}\right)\right| \\
&= \beta^{-k}|t-(\phi_{J|_k}(0)-\phi_{J^*|_k}(0))| = \beta^{-k}|\psi_{J^*|_k}(t)-\phi_{J|_k}(0)| \\
&< \frac{1-N\beta}{\beta(N-1)}.
\end{aligned}$$

Case II. $t_k = N-1$.

In this case, $(j_k, j^*_k) = (N-1, 0)$. This together with the requirements in (B) imply that $\phi_{J|_k}(0)-\psi_{J^*|_k}(t) < \beta^{k-1}(1-\beta)/(N-1)-\beta^k$. By a similar argument as in Case I, we have

$$\sum_{\ell=1}^{\infty}\frac{t_{k+\ell}\beta^{\ell-1}(1-\beta)}{N-1} = \beta^{-k}(\psi_{J^*|_k}(t)-\phi_{J|_k}(0)) > -\frac{1-N\beta}{\beta(N-1)},$$

leading to (5).

The final case $t_k = 1-N$ can be done in the same way as above.

(C) $\Rightarrow$ (B). We will prove by induction that for any $k\geq 1$ and $J\in\Omega_N^k$

$$\mathcal{N}_t(J) = \begin{cases} \{\psi_{J-(t_\ell)_{\ell=1}^k}([t,1+t])\}, & \text{if } J\in\prod_{\ell=1}^k\left(\Omega_N\cap(\Omega_N+t_\ell)\right) \\ \emptyset, & \text{otherwise.}\end{cases} \tag{6}$$

For $k=1$, let $J\in\Omega_N\cap(\Omega_N+t_1)$. In view of the proof of $(B)\Rightarrow(C)$, (5) becomes

$$\begin{cases} \psi_{J-t_1}(t)-\phi_J(0) < (1-\beta)/(N-1)-\beta, & \text{if } t_1 < N-1 \\ \phi_J(0)-\psi_{J-t_1}(t) < (1-\beta)/(N-1)-\beta, & \text{if } t_1 > 1-N.\end{cases}$$

This implies (6) from the geometrical structure of $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$.

Suppose that (6) is true for $k=n$. Let $J=(j_\ell)_{\ell=1}^{n+1}\in\Omega_N^{n+1}$. Then $\mathcal{N}_t(J)=\emptyset$ if $J|_n\notin\prod_{\ell=1}^n\left(\Omega_N\cap(\Omega_N+t_\ell)\right)$. Thus we assume $J|_n\in\prod_{\ell=1}^n\left(\Omega_N\cap(\Omega_N+t_\ell)\right)$. For $j_{n+1}\in\Omega_N\cap(\Omega_N+t_{n+1})$, (5) becomes

$$\begin{cases} \psi_{J-(t_\ell)_{\ell=1}^{n+1}}(t)-\phi_J(0) < \beta^n(1-\beta)/(N-1)-\beta^{n+1}, & \text{if } t_{n+1} < N-1 \\ \phi_J(0)-\psi_{J-(t_\ell)_{\ell=1}^{n+1}}(t) < \beta^n(1-\beta)/(N-1)-\beta^{n+1}, & \text{if } t_{n+1} > 1-N,\end{cases}$$

which implies (6) for $k=n+1$. □

3. The Self-similar structure of $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$

Let Ω be a nonempty finite subset of $\mathbb{Z}$. An infinite string $K\in\Omega^\infty$ is called *strongly periodic with period* q (or simply, strongly periodic) if there exist two finite strings $I=(i_\ell)_{\ell=1}^q, J=(j_\ell)_{\ell=1}^q\in\Omega^q$ with $q\geq 1$ such that $K=I\,J^\infty$ and $I\preccurlyeq J$, where $I\preccurlyeq J$ means $i_\ell\leq j_\ell, 1\leq\ell\leq q$. For two infinite strings $I,J\in\Omega^\infty$, we say $I\preccurlyeq J$ if $I|_k\preccurlyeq J|_k$ for all $k\in\mathbb{N}$. The following lemma (cf. [15, Lemma 3.1]) gives a description of strongly periodic infinite strings.

Lemma 3.1. *Let $(j_\ell)_{\ell=1}^\infty\in\Omega_N^\infty$. If there exists a positive integer q such that $j_{\ell+q}\geq j_\ell$ for all $\ell\in\mathbb{N}$, then $(j_\ell)_{\ell=1}^\infty$ is strongly periodic with period q.*

When t has a unique $\Omega_{\pm N}$-code $(t_\ell)_{\ell=1}^\infty$, from the proof of Theorem 2.2 it follows that there exists a sequence of consecutive subsets $\Omega_N\cap(\Omega_N+t_\ell)$ such that

$$\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)=\pi_N\left(\prod_{\ell=1}^\infty\Omega_N\cap(\Omega_N+t_\ell)\right).$$

Let γ_* be the smallest member of $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$. It is easy to check that

$$\Gamma_t:=\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)-\gamma_*=\pi_N\left(\prod_{\ell=1}^\infty\{0,\ldots,N-1-|t_\ell|\}\right). \tag{7}$$

Thus the Hausdorff and packing dimensions of $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$ are given by (cf. [14])

$$\begin{aligned}\dim_H\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)\quad&=\dim_H\Gamma_t=-\frac{1}{\log\beta}\underline{\lim}_{k\to\infty}\frac{\sum_{\ell=1}^k(N-|t_\ell|)}{k};\\ \dim_P\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)\quad&=\dim_P\Gamma_t=-\frac{1}{\log\beta}\overline{\lim}_{k\to\infty}\frac{\sum_{\ell=1}^k(N-|t_\ell|)}{k}.\end{aligned}$$

The following properties make it easier to deal with Γ_t.

(P5) For $I,J\in\Omega_N^\infty$, if $I\preccurlyeq J$ and $\pi_N(J)\in\Gamma_t$, then $\pi_N(I)\in\Gamma_t$;

(P6) $\Gamma_t=\gamma^*-\Gamma_t$ where $\gamma^*=\pi_N\big((N-1-|t_\ell|)_{\ell=1}^\infty\big)$ is the largest member in Γ_t.

Thus, when Γ_t is generated by an IFS, say $\{f_i(x)=r_ix+b_i\}_{i=1}^p$, we can require all $r_i>0$: if $r_i<0$ we can replace $f_i(x)$ by $f_i^*(x)=-r_ix+b_i+r_i\gamma^*$. This follows from a simple computation (cf. [3, 15])

$$f_i^*(\Gamma_t)=-r_i\Gamma_t+b_i+r_i\gamma^*=r_i(\gamma^*-\Gamma_t)+b_i=r_i\Gamma_t+b_i=f_i(\Gamma_t).$$

Furthermore, we can assume $0=b_1\leq b_2\leq\cdots\leq b_p$ since $0=\pi_N(0^\infty)\in\Gamma_t$ by (P5).

The following theorem gives a sufficient and necessary condition for $t\in\mathcal{S}_{\beta,\pm N}$, i.e., the set of $t\in[-1,1]$ which have a unique $\Omega_{\pm N}$-code and at the same time make the intersection $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$ a self-similar set.

Theorem 3.2. *Given $N\geq 2$ and $\beta\in(1/(2N-1),1/N)$, let $(t_\ell)_{\ell=1}^\infty$ be the unique $\Omega_{\pm N}$-code of $t\in\mathcal{U}_{\beta,\pm N}$. Then $t\in\mathcal{S}_{\beta,\pm N}$ if and only if $(N-1-|t_\ell|)_{\ell=1}^\infty$ is strongly periodic.*

Proof. It suffices to prove that Γ_t, given by (7), is a self-similar set if and only if $(N-1-|t_\ell|)_{\ell=1}^\infty$ is strongly periodic. Firstly, we prove the sufficiency. If $(N-1-|t_\ell|)_{\ell=1}^\infty \in \Omega_N^\infty$ is strongly periodic, it can be written as $(N-1-|t_\ell|)_{\ell=1}^\infty = \sigma\,(\sigma+\tau)^\infty \in \Omega_N^\infty$ where $\sigma = (\sigma_\ell)_{\ell=1}^q, \tau = (\tau_\ell)_{\ell=1}^q \in \Omega_N^q$ for some $q \in \mathbb{N}$ and $\sigma + \tau = (\sigma_\ell + \tau_\ell)_{\ell=1}^q \in \Omega_N^q$. Let

$$\mathcal{S} := \left\{ \beta^{-q} \sum_{\ell=1}^{2q} \frac{j_\ell \beta^{\ell-1}(1-\beta)}{N-1} : \Omega_N^{2q} \ni (j_\ell)_{\ell=1}^{2q} \preccurlyeq \sigma\tau \right\}.$$

One can check that Γ_t can be generated by the IFS $\{f_s(x) = \beta^q(x+s) : s \in \mathcal{S}\}$ (cf. [15]).

Next, we will prove the necessity. By (P6), we can assume that Γ_t is generated by an IFS $\{f_i(x) = r_i x + b_i\}_{i=1}^p$ with $r_i \in (0,1)$ and $0 = b_1 \le b_2 \le \cdots \le b_p$. Note that the union $(0,1) = \bigcup_{q=0}^\infty [\beta^{q+1}, \beta^q)$ is disjoint, there exist some $q \ge 0$ such that $r_1 \in [\beta^{q+1}, \beta^q)$.

Case I. $r_1 = \beta^{q+1}$. Then for each $\ell \ge 1$, it follows from (P5) that

$$\frac{(N-1-|t_\ell|)\beta^{\ell-1}(1-\beta)}{N-1} = \pi_N\big(0^{\ell-1}(N-1-|t_\ell|)0^\infty\big) \in \Gamma_t.$$

Thus

$$f_1\left(\frac{(N-1-|t_\ell|)\beta^{\ell-1}(1-\beta)}{N-1}\right) = \frac{(N-1-|t_\ell|)\beta^{\ell+q}(1-\beta)}{N-1} \in \Gamma_t$$

which implies that $N-1-|t_\ell| \le N-1-|t_{\ell+q+1}|$ for each $\ell \ge 1$. So $(N-1-|t_\ell|)_{\ell=1}^\infty$ is strongly periodic with period $q+1$ by Lemma 3.1.

Case II. $\beta^{q+1} < r_1 < \beta^q$. Let $r_1 = \beta^{q+\gamma}$ with $0 < \gamma < 1$.

(IIa) γ is rational. Take $k \in \mathbb{N}$ such that $k\gamma \in \mathbb{N}$. Note that the IFS $\{f_0(x) = r_1^k x, f_i(x) = r_i x + b_i, 1 \le i \le p\}$ generates Γ_t. Thus the conclusion can be proved in the same way as that in Case I.

(IIb) γ is irrational. Take $k \in \mathbb{N}$ such that

$$\beta < \beta^{1-k\gamma+[k\gamma]} < \frac{1-\beta}{N-1}. \tag{8}$$

This is possible since the set $\{k\gamma - [k\gamma] : k \in \mathbb{N}\}$ is dense in the interval $(0,1)$. Let $f_0(x) = r_1^k x$. Then for some $\beta^{\ell-1}(1-\beta)/(N-1) \in \Gamma_t$ we have

$$f_0\left(\frac{\beta^{\ell-1}(1-\beta)}{N-1}\right) = \frac{\beta^{kq+k\gamma+\ell-1}(1-\beta)}{N-1} < \xi := \frac{\beta^{kq+[k\gamma]+\ell-1}(1-\beta)}{N-1}.$$


$kq + [k\gamma] + \ell$ Level
0
η
ζ
. . .
$kq + [k\gamma] + \ell + 1$ Level
. . .

FIGURE 4. $\xi = (\beta^{kq+[k\gamma]+\ell-1}(1-\beta))/(N-1)$, $\eta = \beta^{kq+[k\gamma]+\ell}$. From the geometrical construction of Γ_t, it is easy to see that $(\eta, \xi) \cap \Gamma_t = \emptyset$.

On the other hand, from (8) it follows that

$$\frac{\beta^{kq+k\gamma+\ell-1}(1-\beta)}{N-1} > \eta := \beta^{kq+[k\gamma]+\ell}.$$

Thus $f_0(\frac{\beta^{\ell-1}(1-\beta)}{N-1}) \notin \Gamma_t$ (see Figure 4), leading to a contradiction. $\square$

In fact, the above proof gives a general result on the structure of a class of subsets of the N-part homogeneous Cantor set.

Corollary 3.3. *Given* $N \geq 2$ *and* $\beta \in (0, 1/N)$*, let* $(i_\ell)_{\ell=1}^\infty, (j_\ell)_{\ell=1}^\infty \in \Omega_N^\infty$ *satisfying* $(i_\ell)_{\ell=1}^\infty \preccurlyeq (j_\ell)_{\ell=1}^\infty$*. Then* $\pi_N(\prod_{\ell=1}^\infty \{i_\ell, i_\ell+1, \ldots, j_\ell\})$ *is a self-similar set if and only if* $(j_\ell - i_\ell)_{\ell=1}^\infty$ *is strongly periodic.*

4. The critical point for $\mathcal{U}_{\beta,\pm N}$

According to a result of Sidorov [19, Proposition 3.8] pertaining to the general digit sets, we have that Lebesgue a.a. $t \in [-1,1]$ have a continuum of distinct $\Omega_{\pm N}$-codes if $\beta \in (1/(2N-1), 1/N)$. However, we will show in this section, for the same set of β's, that there are infinitely many $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code. Note that these t form exactly the set $\mathcal{U}_{\beta,\pm N}$ defined earlier. Moreover, there is a critical point $\beta_c \in (1/(2N-1), 1/N)$ such that $\mathcal{U}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \beta_c)$, and contains countably infinite many elements if $\beta \in (\beta_c, 1/N)$. This can be seen in Theorem 4.6 which is proved by using techniques from beta-expansions.

Given $m \geq 2$ and $\beta \in (1/m, 1)$, let $\Omega_m := \{0, 1, \ldots, m-1\}$. Recall that the sequence $(s_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$ is called a *β-expansion of x with digit set Ω_m* if we can write $x = \sum_{\ell=1}^\infty s_\ell \beta^\ell$ with $s_\ell \in \Omega_m$. The largest number we can obtain in this way is $x_{\max} := (m-1)\beta/(1-\beta)$. Now for any $x \in (0, x_{\max}]$, let us define a sequence $(s_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$ recursively by the *quasi-greedy algorithm* (cf. [20]): let $s_0 = 0$, and if s_ℓ is already defined for all $\ell < n$, then let s_n be the largest element in Ω_m satisfying $\sum_{\ell=1}^n s_\ell \beta^\ell < x$. Obviously, $\sum_{\ell=1}^\infty s_\ell \beta^\ell = x$, and we call $(s_\ell)_{\ell=1}^\infty$ the *quasi-greedy β-expansion of x with digit set Ω_m*. We always call $(s_\ell)_{\ell=1}^\infty$ a quasi-greedy expansion of x if there is no confusion about β and the digit set Ω_m. It is easy to see that $(s_\ell)_{\ell=1}^\infty$ is an *infinite expansion* (i.e., infinitely many s_ℓ are non-zeros).

We use systematically the lexicographical order between sequences: we write $(a_\ell)_{\ell=1}^\infty < (b_\ell)_{\ell=1}^\infty$ or $(b_\ell)_{\ell=1}^\infty > (a_\ell)_{\ell=1}^\infty$ if there exists an $n \in \mathbb{N}$ such that $a_\ell = b_\ell$ for $\ell < n$ and $a_n < b_n$. Furthermore, we write $(a_\ell)_{\ell=1}^\infty \leq (b_\ell)_{\ell=1}^\infty$ or $(b_\ell)_{\ell=1}^\infty \geq (a_\ell)_{\ell=1}^\infty$ if we also allow the equality of the two sequences. Similarly, for two s-blocks $c_1 \ldots c_s$ and $d_1 \ldots d_s$, we write $(c_\ell)_{\ell=1}^s < (d_\ell)_{\ell=1}^s$ if there exists $1 \leq n \leq s$ such that $c_1 \ldots c_{n-1} = d_1 \ldots d_{n-1}$ and $c_n < d_n$. Moreover, we write $(c_\ell)_{\ell=1}^s \leq (d_\ell)_{\ell=1}^s$ if we allow the equality of the two blocks.

Therefore, the quasi-greedy expansion of $x \in (0, x_{\max}]$ is the largest infinite expansion among all the β-expansions of x in the sense of lexicographical order. Note that $1 \in (0, x_{\max}]$ since $\beta > 1/m$. In the remainder of the paper we will reserve the notation $(\delta_\ell)_{\ell=1}^\infty = (\delta_\ell(\beta))_{\ell=1}^\infty$ for the quasi-greedy β-expansion of 1 with digit set Ω_m. The following important properties of the quasi-greedy expansion of 1, will be used in the proof of Theorem 4.6.

Proposition 4.1 (Parry [17]). *Given* $m \geq 2$*, the map* $\beta \to (\delta_\ell(\beta))_{\ell=1}^\infty \in \Omega_m^\infty$*, with* $\beta \in (1/m, 1)$*, is strictly decreasing in the sense of lexicographical order. Moreover, the map is continuous w.r.t. the topology in* Ω_m^∞ *induced by the metric* $d\big((a_\ell)_{\ell=1}^\infty, (b_\ell)_{\ell=1}^\infty\big) = 2^{-\min\{j : a_j \neq b_j\}}$.

Proposition 4.2 (de Vries and Komornik [20])**.** *Given $m \geq 2$ and $\beta \in (1/m, 1)$, let $(\gamma_\ell)_{\ell=1}^\infty$ be an infinite β-expansion of 1 with digit set Ω_m. Then $(\gamma_\ell)_{\ell=1}^\infty$ is the quasi-greedy expansion of 1 if and only if for all $k \geq 1$*

$$\gamma_{k+1}\gamma_{k+2}\cdots \leq \gamma_1\gamma_2\ldots \tag{9}$$

in the lexicographical order.

Given $m \geq 2$, let $\overline{d} = m-1-d$ be the *reflection of the digit* $d \in \Omega_m$. For a sequence $(a_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$, let $\overline{(a_\ell)_{\ell=1}^\infty} = (\overline{a_\ell})_{\ell=1}^\infty = (m-1-a_\ell)_{\ell=1}^\infty$ be the *reflection of the sequence* $(a_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$. A sequence $(a_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$ is said to be *admissible* if for all $k \geq 1$

$$\begin{cases} a_{k+1}a_{k+2}\cdots < a_1a_2\ldots, & \text{if} \quad a_k < m-1 \\ \overline{a_{k+1}a_{k+2}\ldots} < a_1a_2\ldots, & \text{if} \quad a_k > 0. \end{cases}$$

Let $(\tau_\ell)_{\ell=0}^\infty \in \Omega_2^\infty$ be the classical Thue-Morse sequence, i.e., $\tau_0 = 0$, and if τ_ℓ is already defined for some $\ell \geq 0$, set $\tau_{2\ell} = \tau_\ell$ and $\tau_{2\ell+1} = \overline{\tau_\ell} = 1 - \tau_\ell$. Then the sequence $(\tau_\ell)_{\ell=0}^\infty$ begins as follows

$$0\ 1101\ 0011\ 0010\ 1101\ 0010\ 1100\ 1101\ 0011\ 0010\ 1100\ldots.$$

We construct a sequence $(\lambda_\ell)_{\ell=1}^\infty = (\lambda_\ell(m))_{\ell=1}^\infty \in \Omega_m^\infty$ for the even and odd numbers m respectively.

$$\begin{array}{lll} \text{(I).} \quad \lambda_\ell = q-1+\tau_\ell \text{ for } \ell \geq 1, & \text{if} & m = 2q \text{ with } q \geq 1; \\ \text{(II).} \quad \lambda_\ell = q+\tau_\ell-\tau_{\ell-1} \text{ for } \ell \geq 1, & \text{if} & m = 2q+1 \text{ with } q \geq 1. \end{array} \tag{10}$$

Komornik and Loreti [12] showed that $(\lambda_\ell)_{\ell=1}^\infty$ is the smallest admissible sequence in Ω_m^∞ in the sense of lexicographical order. Moreover, they gave the following proposition.

Proposition 4.3 (Komornik and Loreti [12])**.** *Let $(\lambda_\ell)_{\ell=1}^\infty \in \Omega_m^\infty$ be defined in* (10). Then for all $k \geq 1$

$$\lambda_{k+1}\lambda_{k+2}\cdots < \lambda_1\lambda_2\ldots, \quad \overline{\lambda_{k+1}\lambda_{k+2}\ldots} < \lambda_1\lambda_2\ldots.$$

For a more general digit set Ω, there also exist some results on the smallest admissible sequence which is related to the Thue-Morse sequence (cf. [1]).

The following important theorem on the set

$$\mathcal{A}_{\beta,m} := \left\{x \in [0, x_{\max}] : \ x = \sum_{\ell=1}^\infty \varepsilon_\ell \beta^\ell, \ \varepsilon_\ell \in \Omega_m \ \text{ has a unique } \beta\text{-expansion}\right\}$$

is due to Parry [17], Erdös et al. [4], Komornik et al. [12] and de Vries et al. [20].

Theorem 4.4. *Given $m \geq 2$ and $\beta \in (1/m, 1)$, let $(\delta_\ell)_{\ell=1}^\infty$ be the quasi-greedy β-expansion of 1 with digit set Ω_m. Then $\sum_{\ell=1}^\infty \varepsilon_\ell\beta^\ell \in \mathcal{A}_{\beta,m}$ if and only if for all $k \geq 1$*

$$\begin{cases} \varepsilon_{k+1}\varepsilon_{k+2}\cdots < \delta_1\delta_2\ldots, & \textit{if} \quad \varepsilon_k < m-1 \\ \overline{\varepsilon_{k+1}\varepsilon_{k+2}\ldots} < \delta_1\delta_2\ldots, & \textit{if} \quad \varepsilon_k > 0. \end{cases}$$

For $m \geq 2$, let $\beta_{c,m}$ be the unique positive solution of the following equation

$$1 = \sum_{\ell=1}^{\infty} \lambda_\ell \beta^\ell, \tag{11}$$

where $(\lambda_\ell)_{\ell=1}^\infty = (\lambda_\ell(m))_{\ell=1}^\infty \in \Omega_m^\infty$ is defined in (10). We remark here that $\beta_{c,m}$ is a transcendental number for all $m \geq 2$ (cf. [12]). For $m = 2$, Glendinning and Sidorov [7] have shown that the critical point for $\mathcal{A}_{\beta,2}$ is $\beta_{c,2}$, i.e., $\mathcal{A}_{\beta,2}$ has positive Hausdorff dimension if $\beta < \beta_{c,2}$ and $\mathcal{A}_{\beta,2}$ contains at most countably many elements if $\beta > \beta_{c,2}$. Their results can be generalized to the even number case, i.e., for an even number $m \geq 2$, the critical point for $\mathcal{A}_{\beta,m}$ is $\beta_{c,m}$. However, it is more intricate to find the critical point for $\mathcal{A}_{\beta,m}$ for an odd number m. Inspired by [7] we show that for an odd number $m \geq 3$, the critical point for $\mathcal{A}_{\beta,m}$ is still $\beta_{c,m}$, the unique positive solution of Equation (11).

Given $N \geq 2$ and $\beta \in (1/(2N-1), 1/N)$, we will find the critical point for $\mathcal{U}_{\beta,\pm N}$, which is the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code.

To make the connection with the theory of beta-expansions we shift $\Omega_{\pm N}$ to the set

$$\Omega_{\pm N} + N - 1 = \{0, 1, \ldots, 2N-2\} = \Omega_{2N-1}.$$

Thus from $[-1,1] = \pi_{\pm N}\big(\Omega_{\pm N}^\infty\big)$ it follows that

$$[0,2] = \pi_{2N-1}\big(\Omega_{2N-1}^\infty\big) = \left\{ \sum_{\ell=1}^{\infty} \frac{\varepsilon_\ell \beta^{\ell-1}(1-\beta)}{N-1} : \varepsilon_\ell \in \{0,1,\ldots,2N-2\} \right\},$$

where $\pi_{2N-1} := \pi_{\Omega_{2N-1}}$ is as in (1). Let

$$\mathcal{U}_{\beta,2N-1} := \left\{ t \in [0,2] : |\pi_{2N-1}^{-1}(t)| = 1 \right\},$$

i.e., the set of $t \in [0,2]$ having a unique Ω_{2N-1}-code. Thus, it is easy to see that

$$\mathcal{U}_{\beta,2N-1} = \mathcal{U}_{\beta,\pm N} + 1.$$

For $\beta \in (1/(2N-1), 1/N)$, note that

$$x \in \mathcal{A}_{\beta,2N-1} \quad \Longleftrightarrow \quad \frac{1-\beta}{\beta(N-1)} x \in \mathcal{U}_{\beta,2N-1}.$$

Thus Theorem 4.4 yields the the following important theorem which could also be shown in a different way by using (5).

Theorem 4.5. *Given $N \geq 2$ and $\beta \in (1/(2N-1), 1/N)$, let $(\delta_\ell)_{\ell=1}^\infty$ be the quasi-greedy β-expansion of 1 with digit set Ω_{2N-1}. Then $(\varepsilon_\ell)_{\ell=1}^\infty \in \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$ if and only if for all $k \geq 1$*

$$\begin{cases} \varepsilon_{k+1}\varepsilon_{k+2}\cdots < \delta_1\delta_2\ldots, & \text{if} \quad \varepsilon_k \in \{0,\ldots,2N-3\} \\ \overline{\varepsilon_{k+1}\varepsilon_{k+2}\ldots} < \delta_1\delta_2\ldots, & \text{if} \quad \varepsilon_k \in \{1,\ldots,2N-2\}, \end{cases} \tag{12}$$

where $\overline{\varepsilon_{k+1}\varepsilon_{k+2}\ldots}$ is the reflection of $\varepsilon_{k+1}\varepsilon_{k+2}\cdots \in \Omega_{2N-1}^\infty$.

Therefore, dealing with the set $\mathcal{U}_{\beta,\pm N}$ is equivalent to dealing with the set of sequences $(\varepsilon_\ell)_{\ell=1}^\infty \in \Omega_{2N-1}^\infty$ which satisfy (12). Substituting $m = 2N-1$ in (10), we get the smallest admissible sequence $(\lambda_\ell)_{\ell=1}^\infty \in \Omega_{2N-1}^\infty$ which starts with

$$N(N-1)(N-2)N \quad (N-2)(N-1)N(N-1) \quad (N-2)(N-1)N(N-2)\dots.$$

It is helpful to give another equivalent definition of the sequence $(\lambda_\ell)_{\ell=1}^\infty \in \Omega_{2N-1}^\infty$ (cf. [12]), i.e.,

$$\begin{aligned} &\lambda_1 = N, \quad \lambda_{2^{n+1}} = \overline{\lambda_{2^n}} + 1 = 2N-1-\lambda_{2^n} \quad \text{for } n = 0,1,\dots, \\ &\lambda_{2^n+\ell} = \overline{\lambda_\ell} = 2N-2-\lambda_\ell \qquad\quad \text{for } 1 \le \ell < 2^n,\ n = 1,2,\dots. \end{aligned} \tag{13}$$

So it is easy to see $\lambda_{2^n} = N$ for $n = 0,2,4,\dots$ and $\lambda_{2^n} = N-1$ for $n = 1,3,5,\dots$.

Theorem 4.6. *Given* $N \ge 2$, $\beta \in (1/(2N-1), 1/N)$*, let* $\mathcal{U}_{\beta,\pm N}$ *be the set of* $t \in [-1,1]$ *having a unique* $\Omega_{\pm N}$*-code and* $\beta_c \in (1/(2N-1), 1/N)$ *be the unique positive solution of Equation* (11) with $(\lambda_\ell)_{\ell=1}^\infty \in \Omega_{2N-1}^\infty$ defined in (13). Then

(1) If $\beta \in (1/(2N-1), \beta_c)$, then $\dim_H \mathcal{U}_{\beta,\pm N} > 0$;
(2) If $\beta = \beta_c$, then $|\mathcal{U}_{\beta_c,\pm N}| = 2^{\aleph_0}$ and $\dim_H \mathcal{U}_{\beta_c,\pm N} = 0$;
(3) If $\beta \in (\beta_c, 1/N)$, then $|\mathcal{U}_{\beta,\pm N}| = \aleph_0$.

Since $\mathcal{U}_{\beta,\pm N} = \mathcal{U}_{\beta,2N-1} - 1$, the critical point of $\mathcal{U}_{\beta,\pm N}$ is equal to the critical point of $\mathcal{U}_{\beta,2N-1}$. Thus we only need to show the corresponding conclusions for the set $\mathcal{U}_{\beta,2N-1}$.

Using Proposition 4.2 and Proposition 4.3, we obtain $(\delta_\ell(\beta_c))_{\ell=1}^\infty = (\lambda_\ell)_{\ell=1}^\infty$, i.e., $(\lambda_\ell)_{\ell=1}^\infty$ is the quasi-greedy β_c-expansion of 1 with digit set Ω_{2N-1}. The proof of Theorem 4.6 will be divided into several lemmas.

Lemma 4.7. $\lambda_k \dots \lambda_{k+2^n-2} < \lambda_1 \dots \lambda_{2^n-1}$ *for any* $n \ge 2$ *and any* $k \in \{2, \dots, 2^n-1\}$*;* $\overline{\lambda_k \dots \lambda_{k+2^n-2}} < \lambda_1 \dots \lambda_{2^n-1}$ *for any* $n \ge 2$ *and any* $k \in \{1, \dots, 2^n-1\}$*.*

Proof. Since for $n = 2$ the lemma is quickly checked, let $n \ge 3$ and $k \in \{2, \dots, 2^n-1\}$. Then by Proposition 4.3 $\lambda_k\lambda_{k+1}\dots < \lambda_1\lambda_2\dots$, which implies $\lambda_k \dots \lambda_{k+2^n-2} \le \lambda_1 \dots \lambda_{2^n-1}$. It is easy to check that $\lambda_k \dots \lambda_{k+2^n-2} < \lambda_1 \dots \lambda_{2^n-1}$ for $k < 7$. For all other k we can write $k = 2^s + 2^p + j$ with $1 \le p < s < n$ and $1 \le j < 2^p$. It follows from [12, Lemma 5.4] that

$$\lambda_k \dots \lambda_{k+2^{p+1}-j} < \lambda_j \dots \lambda_{2^{p+1}} \le \lambda_1 \dots \lambda_{2^{p+1}-j+1}$$

which implies $\lambda_k \dots \lambda_{k+2^n-2} < \lambda_1 \dots \lambda_{2^n-1}$, since $n > p+1$.

For the second inequality, ignoring the trivial cases $k = 1$ and 2, suppose $k = 2^q + j$ with $1 \le j < 2^q$ and $1 \le q < n$. Then it again follows from [12, Lemma 5.5] that

$$\overline{\lambda_k \dots \lambda_{k+2^q-j}} < \lambda_j \dots \lambda_{2^q} \le \lambda_1 \dots \lambda_{2^q-j+1}.$$

which implies that $\overline{\lambda_k \dots \lambda_{k+2^n-2}} < \lambda_1 \dots \lambda_{2^n-1}$, since $n > q$. ☐

Lemma 4.8. *Let* $n \ge 3$ *be an odd integer. If* $\overline{\lambda_k \dots \lambda_{2^n-1}} = \lambda_1 \dots \lambda_{2^n-k}$ *for some* $k \in \{1, \dots, 2^n-1\}$*, then* $\lambda_{2^n-k+1} = N$*.*

Proof. Suppose $\overline{\lambda_k \dots \lambda_{2^n-1}} = \lambda_1 \dots \lambda_{2^n-k}$. It can not happen that $k < 2^{n-1}$ since then we will obtain that $\overline{\lambda_k \dots \lambda_{k+2^{n-1}-2}} = \lambda_1 \dots \lambda_{2^{n-1}-1}$ which contradicts Lemma 4.7. It is also impossible that $k = 2^{n-1}$ since then $N-2 = \overline{\lambda_{2^{n-1}}} = \lambda_1 = N$. Thus we must have $k > 2^{n-1}$. From the definition of $(\lambda_\ell)_{\ell=0}^\infty$ in (13) it follows that

$$\lambda_{k-2^{n-1}} \dots \lambda_{2^{n-1}-1} = \overline{\lambda_k \dots \lambda_{2^n-1}} = \lambda_1 \dots \lambda_{2^n-k},$$

which implies $N \geq \lambda_{2^n-k+1} \geq \lambda_{2^{n-1}} = N$ by Proposition 4.3. ☐

We want to approximate $(\lambda_\ell)_{\ell=1}^\infty$ by eventually periodic sequences which satisfy (9). This does not work for the obvious choice $(\lambda_1 \dots \lambda_{2^n})^\infty$. Thus we define for $n \geq 0$

$$C_n^\infty = \lambda_1 \dots \lambda_{2^n}(\lambda_{2^n+1} \dots \lambda_{2^{n+1}})^\infty.$$

Since for all $n \geq 0$ we have $\lambda_{2^{n+1}} > \overline{\lambda_{2^n}}$, we obtain that

$$\lambda_1 \dots \lambda_{2^n}(\lambda_{2^n+1} \dots \lambda_{2^{n+1}})^3 > \lambda_1 \dots \lambda_{2^{n+1}} \lambda_{2^{n+1}+1} \dots \lambda_{2^{n+2}},$$

which implies

(P7) $C_0^\infty > C_1^\infty > \cdots > C_n^\infty > \cdots > (\lambda_\ell)_{\ell=1}^\infty$ in the lexicographical order.

Lemma 4.9. *Let $n \geq 3$ be an odd number. Then for any $k \geq 1$ we have $\sigma^k(C_n^\infty) < C_n^\infty$, where σ is the left-shift map.*

Proof. Since C_n^∞ is an eventually periodic sequence in Ω_{2N-1}^∞, we only have to check the lemma for $k \in \{1, \dots, 2^{n+1}-1\}$. For $k = 2^n - 1$ or $2^{n+1}-1$, it is easy to check that $\sigma^k(C_n^\infty) < C_n^\infty$. Then we only need to consider the following two cases.

(I) $k \in \{1, \dots, 2^n - 2\}$. It follows from Lemma 4.7 that

$$\sigma^k(C_n^\infty) = \lambda_{k+1} \dots \lambda_{2^n+k-1} \cdots < \lambda_1 \dots \lambda_{2^n-1}\lambda_{2^n}(\lambda_{2^n+1} \dots \lambda_{2^{n+1}})^\infty = C_n^\infty.$$

(II) $k \in \{2^n, \dots, 2^{n+1}-2\}$. Write $k = 2^n + \ell$. Then, by the definition of $(\lambda_\ell)_{\ell=1}^\infty$,

$$\begin{aligned} \sigma^k(C_n^\infty) &= \lambda_{k+1} \dots \lambda_{2^{n+1}-1}\lambda_{2^{n+1}}(\lambda_{2^n+1} \dots \lambda_{2^{n+1}})^\infty \\ &= \overline{\lambda_{\ell+1} \dots \lambda_{2^n-1}}\lambda_{2^{n+1}}(\lambda_{2^n+1} \dots \lambda_{2^{n+1}})^\infty. \end{aligned}$$

If $\overline{\lambda_{\ell+1} \dots \lambda_{2^n-1}} < \lambda_1 \dots \lambda_{2^n-\ell-1}$, we have shown that $\sigma^k(C_n^\infty) < C_n^\infty$. Otherwise, $\ell \geq 2$ and we have by Proposition 4.3 that $\overline{\lambda_{\ell+1} \dots \lambda_{2^n-1}} = \lambda_1 \dots \lambda_{2^n-\ell-1}$. Using Lemma 4.8 we obtain that also $\lambda_{2^{n+1}} = N = \lambda_{2^n-\ell}$. Thus it is enough to show

$$\lambda_{2^n+1} \dots \lambda_{2^{n+1}-1} < \lambda_{2^n-\ell+1} \dots \lambda_{2^{n+1}-\ell-1}.$$

Taking reflections on both sides, this is equivalent to showing $\lambda_1 \dots \lambda_{2^n-1} > \overline{\lambda_{2^n-\ell+1} \dots \lambda_{2^{n+1}-\ell-1}}$, which is true by Lemma 4.7 since $\ell \geq 2$. ☐

Lemma 4.10. *Let $n \geq 3$ be an odd integer and $\xi_n = (N-1)\lambda_1 \dots \lambda_{2^n-1}, \eta_n = (N-2)\lambda_1 \dots \lambda_{2^n-1}$. Then for any $k \in \{0, \dots, 2^n-1\}$*

$$\begin{aligned} &\sigma^k(\xi_n\eta_n) < \lambda_1 \dots \lambda_{2^{n+1}-k}, \quad \sigma^k(\overline{\xi_n\eta_n}) < \lambda_1 \dots \lambda_{2^{n+1}-k}, \quad \sigma^k(\eta_n\overline{\xi_n}) \leq \lambda_1 \dots \lambda_{2^{n+1}-k}, \\ &\sigma^k(\overline{\eta_n}\xi_n) < \lambda_1 \dots \lambda_{2^{n+1}-k}, \quad \sigma^k(\xi_n\overline{\xi_n}) \leq \lambda_1 \dots \lambda_{2^{n+1}-k}, \quad \sigma^k(\overline{\xi_n}\xi_n) < \lambda_1 \dots \lambda_{2^{n+1}-k}. \end{aligned}$$

Proof. Since the lemma is quickly checked for $k=0$ and 1, we can assume $k \in \{2, \ldots, 2^n - 1\}$. It follows by $\lambda_{2^n} = N-1$ (since n is odd) that

$$\sigma^k(\xi_n\eta_n) = \lambda_k \ldots \lambda_{2^n-1}(N-2)\lambda_1 \ldots \lambda_{2^n-1} < \lambda_k \ldots \lambda_{2^n-1}\lambda_{2^n} \ldots \lambda_{2^{n+1}-1} \leq \lambda_1 \ldots \lambda_{2^{n+1}-k}.$$

For the second inequality, note that $\sigma^k(\overline{\xi_n\eta_n}) = \overline{\lambda_k \ldots \lambda_{2^n-1}} N \overline{\lambda_1 \ldots \lambda_{2^n-1}}$. If $\overline{\lambda_k \ldots \lambda_{2^n-1}} < \lambda_1 \ldots \lambda_{2^n-k}$, we have shown $\sigma^k(\overline{\xi_n\eta_n}) < \lambda_1 \ldots \lambda_{2^{n+1}-k}$. Otherwise, it follows by Proposition 4.3 that $\overline{\lambda_k \ldots \lambda_{2^n-1}} = \lambda_1 \ldots \lambda_{2^n-k}$ which implies $k > 2$. Thus we obtain by Lemma 4.8 that $\lambda_{2^n-k+1} = N$. Hence we only have to show $\overline{\lambda_1 \ldots \lambda_{2^n-1}} < \lambda_{2^n-k+2} \ldots \lambda_{2^{n+1}-k}$ which is equivalent to showing $\lambda_1 \ldots \lambda_{2^n-1} > \overline{\lambda_{2^n-k+2} \ldots \lambda_{2^{n+1}-k}}$. This is true by Lemma 4.7 since $k > 2$. Therefore, $\sigma^k(\overline{\xi_n\eta_n}) < \lambda_1 \ldots \lambda_{2^{n+1}-k}$ for $k \in \{2, \ldots, 2^n - 1\}$. The remaining four inequalities follow from Lemma 4.7 and the fact that for $k \in \{2, \ldots, 2^n - 1\}$

$$\begin{aligned}
\sigma^k(\eta_n\overline{\xi_n}) = \sigma^k(\xi_n\overline{\xi_n}) &= \lambda_k \ldots \lambda_{2^n-1}\overline{(N-1)\lambda_1 \ldots \lambda_{2^n-1}} = \lambda_k \ldots \lambda_{2^{n+1}-1}, \\
\sigma^k(\overline{\eta_n}\xi_n) = \sigma^k(\overline{\xi_n}\xi_n) &= \overline{\lambda_k \ldots \lambda_{2^n-1}}(N-1)\lambda_1 \ldots \lambda_{2^n-1} = \overline{\lambda_k \ldots \lambda_{2^{n+1}-1}}.
\end{aligned}$$

□

From Lemma 4.9 and Proposition 4.2 it follows that C_n^∞ is the quasi-greedy expansion of 1 for some base β_n, i.e., $(\delta_\ell(\beta_n))_{\ell=1}^\infty = C_n^\infty$. Then we obtain from (P7) and Proposition 4.1 that β_n increases to β_c as $n \to \infty$. Thus for $\beta < \beta_c$ there exists a large odd number $n \geq 3$ such that $\beta < \beta_n < \beta_c$, which together with Proposition 4.1 imply that

$$(\delta_\ell(\beta))_{\ell=1}^\infty > (\delta_\ell(\beta_n))_{\ell=1}^\infty = C_n^\infty = \lambda_1 \ldots \lambda_{2^n}(\lambda_{2^n+1} \ldots \lambda_{2^{n+1}})^\infty.$$

It follows from Lemma 4.10 and Theorem 4.5 that

$$X_A^{(n)} \subseteq \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}),$$

where $X_A^{(n)}$ is a subshift of finite type $X_A^{(n)} := \left\{(e_\ell)_{\ell=1}^\infty \in \mathfrak{A}^\infty : \ A(e_\ell, e_{\ell+1}) = 1\right\}$ over the alphabet $\mathfrak{A} = \{\xi_n, \eta_n, \overline{\xi_n}, \overline{\eta_n}\}$ defined by the matrix

$$A = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

It is easy to obtain that $r(A)$, the spectral radius of A, equals $\frac{1+\sqrt{5}}{2}$. Since $\pi_{2N-1}(X_A^{(n)})$ is a graph-directed set satisfying the OSC for large n, we conclude from [16] that

$$\dim_H \mathcal{U}_{\beta,2N-1} \geq \dim_H \pi_{2N-1}(X_A^{(n)}) = \frac{\log r(A)}{-2^n \log \beta} = \frac{\log \frac{1+\sqrt{5}}{2}}{-2^n \log \beta} > 0,$$

which establishes Part (1) of Theorem 4.6.

In the following we will show Part (2) and (3) simultaneously. Let

$$w_n := \lambda_1 \ldots \lambda_{2^n}.$$

Then by the definition of $(\lambda_\ell)_{\ell=1}^\infty$ in (13) it is easy to check that $w_n\overline{w_n} < w_{n+1}$, which implies

(P8) $(w_0\overline{w_0})^\infty < (w_1\overline{w_1})^\infty < \cdots < (w_n\overline{w_n})^\infty < \cdots < (\lambda_\ell)_{\ell=1}^\infty$ in the lexicographical order.

Lemma 4.11. *Given* $N \geq 2$, $\beta \geq \beta_c$ *and* $(\varepsilon_\ell)_{\ell=1}^\infty \in \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$, *if* $\varepsilon_k < 2N-2$ *and* $\varepsilon_{k+1}\cdots\varepsilon_{k+2^n} = w_n$ *for some* $k, n \geq 0$, *then* $\varepsilon_{k+1}\ldots\varepsilon_{k+2^{n+1}} = w_n\overline{w_n}$ *or* w_{n+1}. *Similarly, if* $\varepsilon_k > 0$ *and* $\varepsilon_{k+1}\cdots\varepsilon_{k+2^n} = \overline{w_n}$ *for some* $k, n \geq 0$, *then* $\varepsilon_{k+1}\ldots\varepsilon_{k+2^{n+1}} = \overline{w_n}w_n$ *or* $\overline{w_{n+1}}$.

Proof. Let $(\delta_\ell)_{\ell=1}^\infty := (\delta_\ell(\beta))_{\ell=1}^\infty$. It follows from $\beta \geq \beta_c$ and Proposition 4.1 that

$$(\delta_\ell)_{\ell=1}^\infty \leq (\delta_\ell(\beta_c))_{\ell=1}^\infty = (\lambda_\ell)_{\ell=1}^\infty.$$

Using (12) and the assumption $\varepsilon_k < 2N-2$, we obtain that $\varepsilon_{k+1}\ldots\varepsilon_{k+2^{n+1}} \leq \delta_1\ldots\delta_{2^{n+1}} \leq \lambda_1\ldots\lambda_{2^{n+1}}$. Note that $\varepsilon_{k+1}\cdots\varepsilon_{k+2^n} = w_n = \lambda_1\ldots\lambda_{2^n}$, then $\varepsilon_{k+2^n+1}\ldots\varepsilon_{k+2^{n+1}} \leq \lambda_{2^n+1}\cdots\lambda_{2^{n+1}}$. On the other hand, from (12) and the fact $\varepsilon_{k+2^n} = \lambda_{2^n} > 0$ it follows that $\overline{\varepsilon_{k+2^n+1}\cdots\varepsilon_{k+2^{n+1}}} \leq \delta_1\ldots\delta_{2^n} \leq \lambda_1\ldots\lambda_{2^n}$. Thus by the definition of $(\lambda_\ell)_{\ell=1}^\infty$ in (13), we obtain

$$\lambda_{2^n+1}\cdots\lambda_{2^{n+1}-1}\lambda_{2^{n+1}} \geq \varepsilon_{k+2^n+1}\ldots\varepsilon_{k+2^{n+1}} \geq \overline{\lambda_1\ldots\lambda_{2^n}} = \lambda_{2^n+1}\ldots\lambda_{2^{n+1}-1}(\lambda_{2^{n+1}}-1),$$

which implies $\varepsilon_{k+1}\cdots\varepsilon_{k+2^{n+1}} = w_n\overline{w_n}$ or w_{n+1}.

The result for $\varepsilon_k > 0$ and $\varepsilon_{k+1}\ldots\varepsilon_{k+2^n} = \overline{\lambda_1\ldots\lambda_{2^n}}$ follows similarly. □

Lemma 4.12. *Let* $N \geq 2$ *and* $\beta \in (\beta_c, 1/N)$. *Then there exists some integer* $n^* = n^*(\beta) \geq 0$ *such that* $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1} \setminus \{0,2\})$ *contains only eventually periodic sequences, either with period* 1 *and period block* $N-1$ *or with period* 2^{n+1} *and period block* $w_n\overline{w_n}$ *for some* $n \leq n^*$.

Proof. For $\beta \in (\beta_c, 1/N)$, let $(\delta_\ell)_{\ell=1}^\infty := (\delta_\ell(\beta))_{\ell=1}^\infty$. The proof will be split into two cases: Case I treats $(\delta_\ell)_{\ell=1}^\infty > (w_0\overline{w_0})^\infty$, and Case II treats $(\delta_\ell)_{\ell=1}^\infty \leq (w_0\overline{w_0})^\infty$.

Fix a sequence $(\varepsilon_\ell)_{\ell=1}^\infty \in \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$. In terms of Theorem 4.5, it is easy to see that $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$ is *reflection invariant*, i.e., it contains $(\varepsilon_\ell)_{\ell=1}^\infty$ if and only if it contains $(\overline{\varepsilon_\ell})_{\ell=1}^\infty = (2N-2-\varepsilon_\ell)_{\ell=1}^\infty$. Note that $\overline{N-1} = N-1$ and that the existence of a period block $\overline{w_n}w_n$ implies the existence of a period block $w_n\overline{w_n}$. So we can assume by reflection that $\varepsilon_1 \in \{0,\ldots,N-1\}$. Ignoring the trivial case $(\varepsilon_\ell)_{\ell=1}^\infty = 0^\infty$, let $j \geq 1$ be the least integer such that $\varepsilon_j > 0$. By Proposition 4.1, it follows from $\beta_c < \beta < 1/N$ that

$$(N-1)^\infty = (\delta_\ell(1/N))_{\ell=1}^\infty < (\delta_\ell)_{\ell=1}^\infty < (\delta_\ell(\beta_c))_{\ell=1}^\infty = (\lambda_\ell)_{\ell=1}^\infty,$$

which together with (12) imply $\varepsilon_j \in \{1,\ldots,N\}$. Moreover, we obtain from this with (12) that

$$\varepsilon_{j+1}\varepsilon_{j+2}\cdots \in \prod_1^\infty\{N-2, N-1, N\}.$$

Case I. $(w_0\overline{w_0})^\infty < (\delta_\ell)_{\ell=1}^\infty < (\lambda_\ell)_{\ell=1}^\infty$.

It then follows from (P8) that there exists an integer $n^* \geq 0$ such that $(w_{n^*}\overline{w_{n^*}})^\infty < (\delta_\ell)_{\ell=1}^\infty \leq (w_{n^*+1}\overline{w_{n^*+1}})^\infty$.

(Ia) $\varepsilon_j \in \{1,\ldots,N-1\}$. One case is that $\varepsilon_{j+1}\varepsilon_{j+2}\cdots = (N-1)^\infty$, otherwise, let first $s \geq j$ be the least integer such that $\varepsilon_{s+1} \in \{N, N-2\} = \{w_0, \overline{w_0}\}$, and then let $p = p(s) \geq 0$ be the largest integer such that $\varepsilon_{s+1}\ldots\varepsilon_{s+2^p} = w_p$ or $\overline{w_p}$. Note that when $s > j$, then $0 < \varepsilon_s = N-1 < 2N-2$ or when $s = j$, then $0 < 1 \leq \varepsilon_s \leq N-1 < 2N-2$. Thus substituting $k = s$ and $n = p$ in Lemma 4.11 we obtain $\varepsilon_{s+1}\ldots\varepsilon_{s+2^{p+1}} \in \{w_p\overline{w_p}, \overline{w_p}w_p, w_{p+1}, \overline{w_{p+1}}\}$.

If $\varepsilon_{s+1}\dots\varepsilon_{s+2^{p+1}} = w_{p+1}$ or $\overline{w_{p+1}}$, substituting $k = s$ and $n = p+1$ in Lemma 4.11, we can determine the next 2^{p+1} terms as above. Otherwise, using that $\varepsilon_{s+2^p} = \lambda_{2^p}$ or $\overline{\lambda_{2^p}}$, and then substituting $k = s + 2^p$ and $n = p$ in Lemma 4.11 we can determine the next 2^p terms. This procedure can be continued.

Note that $\varepsilon_{s+1}\varepsilon_{s+2}\dots$ can not have block w_{n^*+1}, otherwise, it follows from (P8) that for some $\ell \geq s$, either

$$\varepsilon_{\ell+1}\varepsilon_{\ell+2}\cdots \geq (w_{n^*+1}\overline{w_{n^*+1}})^\infty \geq (\delta_\ell)_{\ell=1}^\infty$$

with $\varepsilon_\ell < N \leq 2N-2$, or

$$\overline{\varepsilon_{\ell+1}\varepsilon_{\ell+2}\dots} \geq (w_{n^*+1}\overline{w_{n^*+1}})^\infty \geq (\delta_\ell)_{\ell=1}^\infty$$

with $\varepsilon_\ell > N-2 \geq 0$. This is in contradiction with (12).

Therefore, $(\varepsilon_\ell)_{\ell=1}^\infty$ must be eventually periodic either with period block $N-1$ or with period block $w_n\overline{w_n}$ for some $n \leq n^*$.

(Ib) $\varepsilon_j = N$. Let $s = j-1$ in (Ia) and then the result follows by the same argument.

Case II. $(N-1)^\infty < (\delta_\ell)_{\ell=1}^\infty \leq (w_0\overline{w_0})^\infty$.

We conclude in this case that $\varepsilon_{j+1}\varepsilon_{j+2}\cdots = (N-1)^\infty$. Otherwise, there exists a $s \geq j$ such that $\varepsilon_{s+1} = w_0$ or $\overline{w_0}$. Thus by the same argument as in Case I, we obtain for some integer $\ell \geq s$ that either $\varepsilon_{\ell+1}\varepsilon_{\ell+2}\cdots \geq (w_0\overline{w_0})^\infty \geq (\delta_\ell)_{\ell=1}^\infty$ with $\varepsilon_\ell < 2N-2$, or $\overline{\varepsilon_{\ell+1}\varepsilon_{\ell+2}\dots} \geq (w_0\overline{w_0})^\infty \geq (\delta_\ell)_{\ell=1}^\infty$ with $\varepsilon_\ell > 0$, leading to a contradiction with (12). □

Lemma 4.12 yields Part (3) of Theorem 4.6 directly. Let $\mathcal{G}$ be the set of sequences in Ω_{2N-1}^∞ which are eventually periodic with period block $N-1$ or $w_n\overline{w_n}$ for some integer $n \geq 0$. Then the set $\mathcal{G}$ is countable. When $\beta = \beta_c$, it follows from Lemma 4.11 and the proof of Lemma 4.12 that $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta_c,2N-1} \setminus \{0,2\}) \setminus \mathcal{G}$ is included in the set of sequences of the form

$$\tau(w_0\overline{w_0})^{k_0}(w_0\overline{w_{i_1'}})^{k_0'}(w_{i_1}\overline{w_{i_1}})^{k_1}(w_{i_1}\overline{w_{i_2'}})^{k_1'}\dots(w_{i_n}\overline{w_{i_n}})^{k_n}(w_{i_n}\overline{w_{i_{n+1}'}})^{k_n'}\dots,$$

where $\tau \in \bigcup_{k=0}^\infty \Omega_{2N-1}^k$, $k_n \in \mathbb{N}\cup\{0\}$, $k_n' \in \{0,1\}$ and $0 < i_1' \leq i_1 < i_2' \leq i_2 < \cdots \leq i_n < i_{n+1}' \leq i_{n+1} < \dots$, together with their reflections. Thus, since the length of the block w_n is growing exponentially, $\dim_H \mathcal{U}_{\beta_c,2N-1} = 0$ (cf. [6, 7]). Note that $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta_c,2N-1})$ contains the set of sequences of the form

$$(w_0\overline{w_0})^{k_0}\dots(w_n\overline{w_n})^{k_n}\dots,\quad k_n \in \mathbb{N},$$

and the fact that $w_n\overline{w_n}$ can not be written as concatenation of two or more blocks of the form $w_\ell\overline{w_\ell}$ with $\ell < n$. Therefore, $|\mathcal{U}_{\beta_c,2N-1}| = 2^{\aleph_0}$ which yields Part (2), and so finishes the proof of Theorem 4.6.

5. The critical point for $\mathcal{S}_{\beta,\pm N}$

In this section we show that there exist infinitely many $t \in \mathcal{S}_{\beta,\pm N}$, i.e., there exist infinitely many $t \in [-1,1]$ having a unique Ω_{2N-1}-code and making the intersection $\Gamma_{\beta,N}\cap(\Gamma_{\beta,N}+t)$ a self-similar set. Moreover, we find the critical point α_c for $\mathcal{S}_{\beta,\pm N}$, i.e., the set $\mathcal{S}_{\beta,\pm N}$ has positive Hausdorff dimension if $\beta \in (1/(2N-1), \alpha_c)$, and contains countably infinite many elements if $\beta \in [\alpha_c, 1/N)$. We are able to prove that α_c is strictly smaller than β_c, the critical point of $\mathcal{U}_{\beta,\pm N}$ which is the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code.

In order to using techniques from beta-expansions, we consider the set $\mathcal{S}_{\beta,2N-1} = \mathcal{S}_{\beta,\pm N} + 1$. Thus it follows from Theorem 3.2 that for $\beta \in (1/(2N-1), 1/N)$,

$$\mathcal{S}_{\beta,2N-1} = \left\{\pi_{2N-1}((\varepsilon_\ell)_{\ell=1}^\infty) \in \mathcal{U}_{\beta,2N-1} : (N-1-|\varepsilon_\ell - N+1|)_{\ell=1}^\infty \text{ is strongly periodic}\right\}.$$

Let Ψ be a map from Ω_{2N-1} to Ω_N defined by

$$\Psi(\varepsilon) = N-1-|\varepsilon - N+1|,$$

then Ψ induces a map on blocks (for $\xi = \xi_1 \ldots \xi_k \in \Omega_{2N-1}^k$ we let $\Psi(\xi) = \Psi(\xi_1)\ldots\Psi(\xi_k)$), and a map $\Psi_\infty : \Omega_{2N-1}^\infty \to \Omega_N^\infty$ given by $\Psi_\infty((\varepsilon_\ell)_{\ell=1}^\infty) = (\Psi(\varepsilon_\ell))_{\ell=1}^\infty$. Then $\mathcal{S}_{\beta,2N-1}$ can be rewritten as

$$\mathcal{S}_{\beta,2N-1} = \mathcal{U}_{\beta,2N-1} \cap \pi_{2N-1}\Big(\bigcup_{\mathbf{c}} \Psi_\infty^{-1}(\mathbf{c})\Big), \tag{14}$$

where the union is taken over all strongly periodic sequences $\mathbf{c} = (c_\ell)_{\ell=1}^\infty \in \Omega_N^\infty$.

Theorem 5.1. *Given $N \geq 2$ and $\beta \in (1/(2N-1), 1/N)$, let $\Gamma_{\beta,N}$ be the N-part homogeneous Cantor set, and $\mathcal{S}_{\beta,\pm N}$ be the set of $t \in [-1,1]$ having a unique $\Omega_{\pm N}$-code and making the intersection $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a self-similar set. Denote $\alpha_c := [N+1-\sqrt{(N-1)(N+3)}]/2$. Then*

(1) If $\beta \in (1/(2N-1), \alpha_c)$, $\dim_H \mathcal{S}_{\beta,\pm N} > 0$;
(2) If $\beta \in [\alpha_c, 1/N)$, $|\mathcal{S}_{\beta,\pm N}| = \aleph_0$.

Since $\mathcal{S}_{\beta,2N-1} = \mathcal{S}_{\beta,\pm N} + 1$, we only need to consider the corresponding conclusions for $\mathcal{S}_{\beta,2N-1}$. A simple computation yields that α_c satisfies the equation

$$1 = N\alpha_c + \sum_{j=2}^\infty (N-1)\alpha_c^j.$$

Then it follows by Proposition 4.2 that $(\delta_\ell(\alpha_c))_{\ell=1}^\infty = N(N-1)^\infty = \lambda_1\lambda_2^\infty$ is the quasi-greedy α_c-expansion of 1. It follows from Proposition 4.1 and

$$(\delta_\ell(\alpha_c))_{\ell=1}^\infty = \lambda_1\lambda_2^\infty > (\lambda_\ell)_{\ell=1}^\infty = (\delta_\ell(\beta_c))_{\ell=1}^\infty$$

that $\alpha_c < \beta_c$. The proof of Theorem 5.1 will be divided into several lemmas.

Lemma 5.2. *Given $N \geq 2$ and $n \in \mathbb{N}$, let α_n be defined by $(\delta_\ell(\alpha_n))_{\ell=1}^\infty = (N(N-1)^{n-1})^\infty$. If $\beta < \alpha_n$, then $\dim_H \mathcal{S}_{\beta,2N-1} > 0$.*

Proof. Let $v_n = N(N-1)^{n-1}$ and $\overline{v_n} = (N-2)(N-1)^{n-1}$ be its reflection. It follows from $\beta < \alpha_n$ and Proposition 4.1 that $(\delta_\ell(\beta))_{\ell=1}^\infty > (\delta_\ell(\alpha_n))_{\ell=1}^\infty = (N(N-1)^{n-1})^\infty$, which implies that for any $k \in \{0,1,\ldots,n-1\}$

$$\sigma^k(v_nv_n) \leq \delta_1(\alpha_n)\ldots\delta_{2n-k}(\alpha_n), \quad \sigma^k(\overline{v_n}v_n) < \delta_1(\alpha_n)\ldots\delta_{2n-k}(\alpha_n),$$
$$\sigma^k(v_n\overline{v_n}) < \delta_1(\alpha_n)\ldots\delta_{2n-k}(\alpha_n), \quad \sigma^k(\overline{v_n}\overline{v_n}) < \delta_1(\alpha_n)\ldots\delta_{2n-k}(\alpha_n).$$

Thus by Theorem 4.5 we obtain that

$$\prod_1^\infty \{v_n, \overline{v_n}\} \subseteq \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}).$$

Since $\Psi(v_n) = (N-2)(N-1)^{n-1} = \Psi(\overline{v_n})$, it is easy to see that

$$\prod_1^\infty \{v_n, \overline{v_n}\} \subseteq \Psi_\infty^{-1}\big(((N-2)(N-1)^{n-1})^\infty\big).$$

Thus noting that $((N-2)(N-1)^{n-1})^\infty$ is obviously a strongly periodic sequence in Ω_N^∞, it follows from (14) that

$$\prod_1^\infty \{v_n, \overline{v_n}\} \subseteq \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}\big(((N-2)(N-1)^{n-1})^\infty\big) \subseteq \pi_{2N-1}^{-1}(\mathcal{S}_{\beta,2N-1})$$

which implies $\dim_H \mathcal{S}_{\beta,2N-1} \geq \dim_H \pi_{2N-1}(\prod_1^\infty \{v_n, \overline{v_n}\}) > 0$. □

Since $(\delta_\ell(\alpha_n))_{\ell=1}^\infty = (N(N-1)^{n-1})^\infty$ decreases to $N(N-1)^\infty = (\delta_\ell(\alpha_c))_{\ell=1}^\infty$ in the sense of lexicographical order as $n \to \infty$, we obtain from Proposition 4.1 that α_n increases to α_c. Thus for each $\beta < \alpha_c$, there exists some $n \in \mathbb{N}$ such that $\beta < \alpha_n$ and then $\dim_H \mathcal{S}_{\beta,2N-1} > 0$ by Lemma 5.2. This finishes the proof of Part (1) of Theorem 5.1.

In the following we will show Part (2). For $\beta \in [\alpha_c, 1/N)$, it follows by Proposition 4.1 that $(\delta_\ell(\beta))_{\ell=1}^\infty \leq (\delta_\ell(\alpha_c))_{\ell=1}^\infty = N(N-1)^\infty$, which together with Theorem 4.5 imply the following property:

(P9) For $N \geq 2$ and $\beta \in [\alpha_c, 1/N)$, any block in $\mathcal{F}$ is forbidden in $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$ where

$$\mathcal{F} = \bigcup_{k=0}^\infty \bigcup_{\tau=N-2}^{N-1} \{\tau N(N-1)^k N,\ \overline{\tau}(N-2)(N-1)^k(N-2)\}.$$

For a positive integer n, let $\mathcal{B}_n$ be the set of blocks of length n occurring in elements of $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$, i.e.,

$$\mathcal{B}_n := \big\{\varepsilon_{i+1}\varepsilon_{i+2}\ldots\varepsilon_{i+n} :\ i \geq 0,\ (\varepsilon_\ell)_{\ell=1}^\infty \in \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})\big\}.$$

Lemma 5.3. *Given $N \geq 2$ and $\beta \in [\alpha_c, 1/N)$, let $\mathbf{b} = b_1 \ldots b_p \in \{N-2, N-1\}^p$ with $b_1 = N-1$ for some $p \in \mathbb{N}$. Then $\Psi^{-1}(\mathbf{b}) \cap \mathcal{B}_p = \{(N-1)^p\}$ or $\{\xi, \overline{\xi}\}$ for some $\xi \in \{N-2, N-1, N\}^p$.*

Proof. Let $\xi = \xi_1 \ldots \xi_p \in \Psi^{-1}(\mathbf{b}) \cap \mathcal{B}_p$. Then it follows from $\mathbf{b} \in \{N-2, N-1\}^p$ and the definition of Ψ that $\xi \in \{N-2, N-1, N\}^p$. Note that $\Psi^{-1}(N-1) = \{N-1\}$ and $\Psi^{-1}(N-2) = \{N-2, N\}$.

(I) $\mathbf{b} = (N-1)^p$. Then $\Psi^{-1}(\mathbf{b}) \cap \mathcal{B}_p = \{(N-1)^p\}$.

(II) $\mathbf{b} \neq (N-1)^p$. Let $b_{k_1} = b_{k_2} = \cdots = b_{k_s} = N-2$ for $1 < k_1 < k_2 < \cdots < k_s \leq p$, and $b_k = N-1$ for $k \neq k_i$. Then also $\xi_k = N-1$ for $k \neq k_i$. Moreover, if $\xi_{k_1} = N$, then it follows from (P9) that $\xi_{k_2} = N-2,\ \xi_{k_3} = N,\ \xi_{k_4} = N-2$ and so on. Similarly, if $\xi_{k_1} = N-2$ we will obtain by (P9) that $\xi_{k_2} = N,\ \xi_{k_3} = N-2,\ \xi_{k_4} = N$ and so on. Thus, $\Psi^{-1}(\mathbf{b}) \cap \mathcal{B}_p = \{\xi, \overline{\xi}\}$. □

Lemma 5.4. *Given $N \geq 2$ and $\beta \in [\alpha_c, 1/N)$, let $\boldsymbol{c} = (c_\ell)_{\ell=1}^\infty \in \Omega_N^\infty$ be a strongly periodic sequence. Then $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\boldsymbol{c})$ is at most countable.*

Proof. Note by $\beta \geq \alpha_c$ that $(\delta_\ell(\beta))_{\ell=1}^\infty \leq (\delta_\ell(\alpha_c))_{\ell=1}^\infty = N(N-1)^\infty$. Thus for any sequence $(\varepsilon_\ell)_{\ell=1}^\infty \in \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$, we obtain by the same argument as in Lemma 4.12 that

$$\varepsilon_k \varepsilon_{k+1} \cdots \in \prod_1^\infty \{N-2, N-1, N\}$$

for some large $k \in \mathbb{N}$, which implies that $\Psi_\infty(\varepsilon_k \varepsilon_{k+1} \ldots) \in \{N-2, N-1\}^\infty$. Let $\mathbf{c} = a_1 \ldots a_q (b_1 \ldots b_q)^\infty$ with $a_\ell \leq b_\ell$, $1 \leq \ell \leq q$ be a strongly periodic sequence in Ω_N^∞ such that $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c}) \neq \emptyset$. Then

$$b_1 \ldots b_q \in \{N-2, N-1\}^q.$$

Case I. $b_1 \ldots b_q = (N-2)^q$. It follows from $\Psi^{-1}(N-2) = \{N-2, N\}$ that $\Psi_\infty^{-1}(\mathbf{c}) \subseteq \Psi^{-1}(a_1 \ldots a_q)\{N-2, N\}^\infty$. Note by (P9) (with $\tau = N-2, k = 0$) that blocks $N(N-2)^2$ and $(N-2)N^2$ are forbidden in $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$. Thus

$$\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c}) \subseteq \Psi^{-1}(a_1 \ldots a_q)\{N^\infty, (N(N-2))^\infty, ((N-2)N)^\infty, (N-2)^\infty\}$$

which is at most countable.

Case II. $b_1 \ldots b_q \neq (N-2)^q$. Then there exists $b_k = N-1$ for some $k \in \{1, \ldots, q\}$. Note that

$$\mathbf{c} = a_1 \ldots a_q (b_1 \ldots b_q)^\infty = a_1 \ldots a_q b_1 \ldots b_{k-1} (b_k \ldots b_q b_1 \ldots b_{k-1})^\infty.$$

It follows from Lemma 5.3 that there exists a q-block $\xi = \xi_1 \ldots \xi_q \in \{N-2, N-1, N\}^q$ such that $\Psi^{-1}(b_k \ldots b_q b_1 \ldots b_{k-1}) \cap \mathcal{B}_q = \{\xi, \overline{\xi}\}$. Thus

$$\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c}) \subseteq \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Big(\Psi^{-1}(a_1 \ldots a_q b_1 \ldots b_{k-1}) \prod_1^\infty \{\xi, \overline{\xi}\}\Big).$$

Note that since $\Psi^{-1}(\mathbf{c})$ and $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1})$ are all reflection invariant, $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c})$ is also reflection invariant. Thus we only need to consider the following three cases.

(IIa) $\xi = (N-1)^q$. Then $\prod_1^\infty \{\xi, \overline{\xi}\}$ collapses to a single point $(N-1)^\infty$.

(IIb) $\xi = (N-1)^\ell N \xi_{\ell+2} \ldots \xi_{q-r-1} N (N-1)^r$ with $\ell \geq 1, r \geq 0$ and $\ell + r \leq q-1$ (note that $\xi = (N-1)^\ell N (N-1)^r$ if $\ell + r = q-1$). It follows by (P9) that blocks $\xi\xi$ and $\overline{\xi\xi}$ are forbidden in $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c})$. Thus $\prod_1^\infty \{\xi, \overline{\xi}\}$ collapses to two points $(\xi\overline{\xi})^\infty$ and $(\overline{\xi}\xi)^\infty$.

(IIc) $\xi = (N-1)^\ell N \xi_{\ell+2} \ldots \xi_{q-r-1} (N-2)(N-1)^r$ with $\ell \geq 1, r \geq 0$ and $\ell + r \leq q-2$. By the same argument as in (IIb) we also obtain that $\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi^{-1}(\mathbf{c})$ is at most countable. □

It follows from Lemma 5.4 and (14) that for $\beta \in [\alpha_c, 1/N)$, the set

$$\pi_{2N-1}^{-1}(\mathcal{S}_{\beta,2N-1}) = \pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \bigcup_{\mathbf{c}} \Psi_\infty^{-1}(\mathbf{c}) = \bigcup_{\mathbf{c}} \big(\pi_{2N-1}^{-1}(\mathcal{U}_{\beta,2N-1}) \cap \Psi_\infty^{-1}(\mathbf{c})\big)$$

is at most countable since the union on the right is countable. Note that for $\beta \in [\alpha_c, 1/N)$, $\{0^q(N-1)^\infty : q \in \mathbb{N}\} \subseteq \pi_{2N-1}^{-1}(\mathcal{S}_{\beta,2N-1})$. This gives Part (2), finishing the proof of Theorem 5.1.

6. Final remarks

In this paper we determined the size of two types of sets $\mathcal{U}_{\beta,\pm N}$, and $\mathcal{S}_{\beta,\pm N}$, where $\mathcal{U}_{\beta,\pm N}$ is the set of $t \in \Gamma_{\beta,N} - \Gamma_{\beta,N}$ having a unique $\Omega_{\pm N}$-code and $\mathcal{S}_{\beta,\pm N}$ is the set of t not only having a unique code but also making the intersection $\Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t)$ a self-similar set. It follows from [19] that for $\beta \in (1/(2N-1), 1/N)$ there also exist a lot of $t \in \Gamma_{\beta,N} - \Gamma_{\beta,N} = [-1,1]$ having exactly p different $\Omega_{\pm N}$-codes for any integer $p \geq 2$. Let

$$\mathcal{F}_{\beta,\pm N}^{(p)} := \{t \in \Gamma_{\beta,N} - \Gamma_{\beta,N} : t \text{ has exactly } p \text{ different } \Omega_{\pm N}\text{-codes}\},$$

and

$$\mathcal{S}_{\beta,\pm N}^{(p)} := \{t \in \mathcal{F}_{\beta,\pm N}^{(p)} : \Gamma_{\beta,N} \cap (\Gamma_{\beta,N} + t) \ \text{ is a self-similar set}\}.$$

Problem. How large is the set $\mathcal{F}_{\beta,\pm N}^{(p)}$ for a given positive integer $p \geq 2$? How to characterize this set? This is also an open problem for beta-expansions. Moreover, how large is the set $\mathcal{S}_{\beta,\pm N}^{(p)}$?

References

[1] Allouche J and Frougny C 2009 Univoque numbers and an avatar of Thue-Morse *Acta Arith.* **136** 319-329

[2] Davis G and Hu T Y 1995 On the structure of the intersection of two middle third Cantor sets *Publ. Mat.* **39** 43-60

[3] Deng G T, He X G and Wen Z X 2008 Self-similar structure on intersection of triadic Cantor sets *J. Math. Anal. Appl.* **337** 617-631

[4] Erdös P, Joó I and Komornik V 1990 Characterization of the unique expansions $1 = \sum_{i=1}^{\infty} q^{-n_i}$ and related problems *Bull. Soc. Math. France* **118** 377-390

[5] Falconer K J 1990 *Fractal Geometry–Mathematical Foundations and Applications* (Chichester: John Wiley & Sons Ltd.)

[6] Fröberg C 1977 Accurate estimation of the number of binary partitions *BIT Numerical Mathematics* **17** 386-391

[7] Glendinning P and Sidorov N 2001 Unique representations of real numbers in non-integer bases *Math. Res. Lett.* **8** 535-543

[8] Kenyon R and Peres Y 1991 Intersecting random translates of invariant Cantor sets *Invent. Math.* **104** 601-629

[9] Kraft R 1992 Intersection of thick Cantor sets *Memoirs of AMS* **97** vi+119 pp

[10] Kraft R 1994 What's the difference between Cantor sets? *The Amer. Math. Monthly* **101** 640-650

[11] Kraft R 1999 Random intersection of thick Cantor sets *Trans. Amer. Math. Soc.* **352** 1315-1328

[12] Komornik V and Loreti P 2002 Subexpansions, supexpansions and uniqueness properties in non-integer bases *Periodica Mathematica Hungarica* **44(2)** 197-218

[13] Li W X and Xiao D M 1998 On the intersection of translation of middle-α Cantor sets *Fractals and Beyond-Complexities in the Sciences (Valletta, 1998)*(Singapore: World Scientific) 137-148

[14] Li W X and Xiao D M 1998 A note on generalized Moran set *Acta Mathematica Scientia* **18**(supp.) 88–93

[15] Li W X, Yao Y Y and Zhang Y X Self-similar structure on intersection of homogeneous symmetric Cantor sets *Mathematische Nachrichten.* at press

[16] Mauldin R and Williams S 1988 Hausdorff dimension in graph directed constructions *Trans. Amer. Math. Soc.* **309(2)** 811–829

[17] Parry W 1960 On the β-expansions of real numbers *Acta Math. Acad. Sci. Hungary* **11** 401-416

[18] Rényi A 1957 Representations for real numbers and their ergodic properties *Acta Math. Hungar.* **8** 477-493

[19] Sidorov N 2007 Combinatorics of linear iterated function systems with overlaps *Nonlinearity* **20** 1299-1312

[20] Vries M and Komornik V 2009 Unique expansions of real numbers *Adv. Math.* **221** 390-427

[21] Zou Y R, Lu J and Li W X 2008 Self-similar structure on the intersection of middle-$(1-2\beta)$ Cantor sets with $\beta \in (1/3, 1/2)$ *Nonlinearity* **21** 2899-2910

3TU APPLIED MATHEMATICS INSTITUTE AND DELFT UNIVERSITY OF TECHNOLOGY, FACULTY EWI, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS.

WENXIA LI. DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, PEOPLES REPUBLIC OF CHINA

E-mail address: `D.Kong@tudelft.nl`, `wxli@math.ecnu.edu.cn`, `F.M.Dekking@tudelft.nl`